

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use LEXIVA safely and effectively. See full prescribing information for LEXIVA.

LEXIVA (fosamprenavir calcium) Tablets and Oral Suspension
Initial U.S. Approval: 2003

INDICATIONS AND USAGE

LEXIVA is an HIV protease inhibitor indicated in combination with other antiretroviral agents for the treatment of HIV-1 infection. (1)

DOSAGE AND ADMINISTRATION

- Therapy-Naive Adults: LEXIVA 1,400 mg twice daily; LEXIVA 1,400 mg once daily plus ritonavir 200 mg once daily; LEXIVA 1,400 mg once daily plus ritonavir 100 mg once daily; LEXIVA 700 mg twice daily plus ritonavir 100 mg twice daily. (2.1)
- Protease Inhibitor-Experienced Adults: LEXIVA 700 mg twice daily plus ritonavir 100 mg twice daily. (2.1)
- Pediatric Patients (2 to 18 years of age): Dosage should be calculated based on body weight (kg) and should not exceed adult dose. (2.2)
- Hepatic Impairment: Recommended adjustments for patients with mild, moderate, or severe hepatic impairment. (2.3)

Dosing Considerations

- LEXIVA Tablets may be taken with or without food. (2)
- LEXIVA Suspension: Adults should take without food; pediatric patients should take with food. (2)

DOSAGE FORMS AND STRENGTHS

700-mg tablets and 50-mg/mL oral suspension (3)

CONTRAINDICATIONS

- Hypersensitivity to LEXIVA or amprenavir (e.g., Stevens-Johnson syndrome). (4)
- Drugs highly dependent on CYP3A4 for clearance and for which elevated plasma levels may result in serious and/or life-threatening events. (4)
- Review ritonavir contraindications when used in combination. (4)

WARNINGS AND PRECAUTIONS

- Certain drugs should not be coadministered with LEXIVA due to risk of serious or life-threatening adverse reactions. (5.1)

- LEXIVA should be discontinued for severe skin reactions including Stevens-Johnson syndrome. (5.2)
- LEXIVA should be used with caution in patients with a known sulfonamide allergy. (5.3)
- Use of higher than approved doses may lead to transaminase elevations. Patients with hepatitis B or C are at increased risk of transaminase elevations. (5.4)
- Patients receiving LEXIVA may develop new onset or exacerbations of diabetes mellitus, hyperglycemia (5.5), immune reconstitution syndrome (5.6), redistribution/accumulation of body fat (5.7), and elevated triglyceride and cholesterol concentrations (5.8). Monitor cholesterol and triglycerides prior to therapy and periodically thereafter.
- Acute hemolytic anemia has been reported with amprenavir. (5.9)
- Hemophilia: Spontaneous bleeding may occur, and additional factor VIII may be required. (5.10)
- Nephrolithiasis: Cases of nephrolithiasis have been reported with fosamprenavir. (5.11)

ADVERSE REACTIONS

- In adults the most common adverse reactions (incidence greater than or equal to 4%) are diarrhea, rash, nausea, vomiting, headache. (6.1)
- Vomiting was more frequent in pediatrics than in adults. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact ViiV Healthcare at 1-877-844-8872 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Coadministration of LEXIVA with drugs that induce CYP3A4 may decrease amprenavir (active metabolite) concentrations leading to potential loss of virologic activity. (7, 12.3)
- Coadministration with drugs that inhibit CYP3A4 may increase amprenavir concentrations. (7, 12.3)
- Coadministration of LEXIVA and ritonavir may result in clinically significant interactions with drugs metabolized by CYP2D6. (7)

See 17 for PATIENT COUNSELING INFORMATION and FDA-Approved Patient Labeling.

Revised: May 2011

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Adults

2.2 Pediatric Patients (Aged 2 to 18 Years)

2.3 Patients With Hepatic Impairment

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Drug Interactions

5.2 Skin Reactions

5.3 Sulfa Allergy

5.4 Hepatic Toxicity

5.5 Diabetes/Hyperglycemia

5.6 Immune Reconstitution Syndrome

5.7 Fat Redistribution

5.8 Lipid Elevations

5.9 Hemolytic Anemia

5.10 Patients With Hemophilia

5.11 Nephrolithiasis

5.12 Resistance/Cross-Resistance

6 ADVERSE REACTIONS

6.1 Clinical Trials

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

7.1 CYP Inhibitors and Inducers

7.2 Drugs That Should Not Be Coadministered With LEXIVA

7.3 Established and Other Potentially Significant Drug Interactions

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.3 Pharmacokinetics

12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

14.1 Therapy-Naive Adult Patients

14.2 Protease Inhibitor-Experienced Adult Patients

14.3 Pediatric Patients

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

17.1 Drug Interactions

17.2 Sulfa Allergy

17.3 Redistribution/Accumulation of Body Fat

17.4 Information About Therapy With LEXIVA

17.5 Oral Suspension

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

LEXIVA[®] is indicated in combination with other antiretroviral agents for the treatment of human immunodeficiency virus (HIV-1) infection.

The following points should be considered when initiating therapy with LEXIVA plus ritonavir in protease inhibitor-experienced patients:

- The protease inhibitor-experienced patient study was not large enough to reach a definitive conclusion that LEXIVA plus ritonavir and lopinavir plus ritonavir are clinically equivalent [see *Clinical Studies (14.2)*].
- Once-daily administration of LEXIVA plus ritonavir is not recommended for adult protease inhibitor-experienced patients or any pediatric patients.

2 DOSAGE AND ADMINISTRATION

LEXIVA Tablets may be taken with or without food.

Adults should take LEXIVA Oral Suspension without food. Pediatric patients should take LEXIVA Oral Suspension with food [see *Clinical Pharmacology (12.3)*]. If emesis occurs within 30 minutes after dosing, re-dosing of LEXIVA Oral Suspension should occur.

Higher-than-approved dose combinations of LEXIVA plus ritonavir are not recommended due to an increased risk of transaminase elevations [see *Overdosage (10)*].

When LEXIVA is used in combination with ritonavir, prescribers should consult the full prescribing information for ritonavir.

2.1 Adults

Therapy-Naive Adults:

- LEXIVA 1,400 mg twice daily (without ritonavir).
- LEXIVA 1,400 mg once daily plus ritonavir 200 mg once daily.
- LEXIVA 1,400 mg once daily plus ritonavir 100 mg once daily.

Dosing of LEXIVA 1,400 mg once daily plus ritonavir 100 mg once daily is supported by pharmacokinetic data [see *Clinical Pharmacology (12.3)*].

- LEXIVA 700 mg twice daily plus ritonavir 100 mg twice daily.

Dosing of LEXIVA 700 mg twice daily plus 100 mg ritonavir twice daily is supported by pharmacokinetic and safety data [see *Clinical Pharmacology (12.3)*].

Protease Inhibitor-Experienced Adults:

- LEXIVA 700 mg twice daily plus ritonavir 100 mg twice daily

2.2 Pediatric Patients (Aged 2 to 18 Years)

The recommended dosage of LEXIVA in patients aged greater than or equal to 2 years should be calculated based on body weight (kg) and should not exceed the recommended adult dose. The data are insufficient to recommend: (1) once-daily dosing of LEXIVA alone or in combination with ritonavir, and (2) any dosing of LEXIVA in therapy-experienced patients aged 2 to 5 years.

Therapy-Naive Aged 2 to 5 Years:

- LEXIVA Oral Suspension 30 mg/kg twice daily, not to exceed the adult dose of LEXIVA 1,400 mg twice daily.

Therapy-Naive Aged Greater Than or Equal to 6 Years:

- Either LEXIVA Oral Suspension 30 mg/kg twice daily not to exceed the adult dose of LEXIVA 1,400 mg twice daily or LEXIVA Oral Suspension 18 mg/kg plus ritonavir 3 mg/kg twice daily not to exceed the adult dose of LEXIVA 700 mg plus ritonavir 100 mg twice daily.

Therapy-Experienced Aged Greater Than or Equal to 6 Years:

- LEXIVA Oral Suspension 18 mg/kg plus ritonavir 3 mg/kg administered twice daily not to exceed the adult dose of LEXIVA 700 mg twice daily plus ritonavir 100 mg twice daily.

Other Dosing Considerations:

- When administered without ritonavir, the adult regimen of LEXIVA Tablets 1,400 mg twice daily may be used for pediatric patients weighing at least 47 kg.
- When administered in combination with ritonavir, LEXIVA Tablets may be used for pediatric patients weighing at least 39 kg; ritonavir capsules may be used for pediatric patients weighing at least 33 kg.

2.3 Patients With Hepatic Impairment

See Clinical Pharmacology (12.3).

Mild Hepatic Impairment (Child-Pugh Score Ranging From 5 to 6): LEXIVA should be used with caution at a reduced dosage of 700 mg twice daily without ritonavir (therapy-naive) or 700 mg twice daily plus ritonavir 100 mg once daily (therapy-naive or protease inhibitor-experienced).

Moderate Hepatic Impairment (Child-Pugh Score Ranging From 7 to 9): LEXIVA should be used with caution at a reduced dosage of 700 mg twice daily without ritonavir (therapy-naive), or 450 mg twice daily plus ritonavir 100 mg once daily (therapy-naive or protease inhibitor-experienced).

Severe Hepatic Impairment (Child-Pugh Score Ranging From 10 to 15): LEXIVA should be used with caution at a reduced dosage of 350 mg twice daily without ritonavir (therapy-naive) or 300 mg twice daily plus ritonavir 100 mg once daily (therapy-naive or protease inhibitor-experienced).

3 DOSAGE FORMS AND STRENGTHS

LEXIVA Tablets, 700 mg, are pink, film-coated, capsule-shaped, biconvex tablets with “GX LL7” debossed on one face.

LEXIVA Oral Suspension, 50 mg/mL, is a white to off-white suspension that has a characteristic grape-bubblegum-peppermint flavor.

4 CONTRAINDICATIONS

LEXIVA is contraindicated:

- in patients with previously demonstrated clinically significant hypersensitivity (e.g., Stevens-Johnson syndrome) to any of the components of this product or to amprenavir.
- when coadministered with drugs that are highly dependent on CYP3A4 for clearance and for which elevated plasma concentrations are associated with serious and/or life-threatening events (Table 1).

Table 1. Drugs Contraindicated With LEXIVA (Information in the table applies to LEXIVA with or without ritonavir, unless otherwise indicated.)

Drug Class/Drug Name	Clinical Comment
Alpha 1-adrenoreceptor antagonist: Alfuzosin	Potentially increased alfuzosin concentrations can result in hypotension.
Antiarrhythmics: Flecainide, propafenone	POTENTIAL for serious and/or life-threatening reactions such as cardiac arrhythmias secondary to increases in plasma concentrations of antiarrhythmics if LEXIVA is co-prescribed with ritonavir .
Antimycobacterials: Rifampin ^a	May lead to loss of virologic response and possible resistance to LEXIVA or to the class of protease inhibitors.
Ergot derivatives: Dihydroergotamine, ergonovine, ergotamine, methylergonovine	POTENTIAL for serious and/or life-threatening reactions such as acute ergot toxicity characterized by peripheral vasospasm and ischemia of the extremities and other tissues.
GI motility agents: Cisapride	POTENTIAL for serious and/or life-threatening reactions such as cardiac arrhythmias.
Herbal products: St. John's wort (<i>hypericum perforatum</i>)	May lead to loss of virologic response and possible resistance to LEXIVA or to the class of protease inhibitors.
HMG co-reductase inhibitors: Lovastatin, simvastatin	POTENTIAL for serious reactions such as risk of myopathy including rhabdomyolysis.
Neuroleptic: Pimozide	POTENTIAL for serious and/or life-threatening reactions such as cardiac arrhythmias.
Non-nucleoside reverse transcriptase inhibitor: Delavirdine ^a	May lead to loss of virologic response and possible resistance to delavirdine.
PDE5 inhibitor: Sildenafil (REVATIO [®]) (for treatment of pulmonary arterial hypertension)	A safe and effective dose has not been established when used with LEXIVA. There is increased potential for sildenafil-associated adverse events (which include visual disturbances, hypotension, prolonged erection, and syncope).
Sedative/hypnotics: Midazolam, triazolam	POTENTIAL for serious and/or life-threatening reactions such as prolonged or increased sedation or respiratory depression.

^a See *Clinical Pharmacology (12.3) Tables 10, 11, 12, or 13 for magnitude of interaction.*

- when coadministered with ritonavir in patients receiving the antiarrhythmic agents flecainide and propafenone. If LEXIVA is coadministered with ritonavir, reference should be made to the full prescribing information for ritonavir for additional contraindications.

5 WARNINGS AND PRECAUTIONS

5.1 Drug Interactions

See Table 1 for listings of drugs that are contraindicated due to potentially life-threatening adverse events, significant drug interactions, or due to loss of virologic activity [see *Contraindications (4)*, *Drug Interactions (7.2)*]. See Table 6 for a listing of established and other potentially significant drug interactions [see *Drug Interactions (7.3)*].

5.2 Skin Reactions

Severe and life-threatening skin reactions, including 1 case of Stevens-Johnson syndrome among 700 patients treated with LEXIVA in clinical studies. Treatment with LEXIVA should be discontinued for severe or life-threatening rashes and for moderate rashes accompanied by systemic symptoms [see *Adverse Reactions (6)*].

5.3 Sulfa Allergy

LEXIVA should be used with caution in patients with a known sulfonamide allergy. Fosamprenavir contains a sulfonamide moiety. The potential for cross-sensitivity between drugs in the sulfonamide class and fosamprenavir is unknown. In a clinical study of LEXIVA used as the sole protease inhibitor, rash occurred in 2 of 10 patients (20%) with a history of sulfonamide allergy compared with 42 of 126 patients (33%) with no history of sulfonamide allergy. In 2 clinical studies of LEXIVA plus low-dose ritonavir, rash occurred in 8 of 50 patients (16%) with a history of sulfonamide allergy compared with 50 of 412 patients (12%) with no history of sulfonamide allergy.

5.4 Hepatic Toxicity

Use of LEXIVA with ritonavir at higher-than-recommended dosages may result in transaminase elevations and should not be used [see *Dosage and Administration (2)*, *Overdosage (10)*]. Patients with underlying hepatitis B or C or marked elevations in transaminases prior to treatment may be at increased risk for developing or worsening of transaminase elevations. Appropriate laboratory testing should be conducted prior to initiating therapy with LEXIVA and patients should be monitored closely during treatment.

5.5 Diabetes/Hyperglycemia

New onset diabetes mellitus, exacerbation of pre-existing diabetes mellitus, and hyperglycemia have been reported during postmarketing surveillance in HIV-infected patients receiving protease inhibitor therapy. Some patients required either initiation or dose adjustments of insulin or oral hypoglycemic agents for treatment of these events. In some cases, diabetic ketoacidosis has occurred. In those patients who discontinued protease inhibitor therapy, hyperglycemia persisted in some cases. Because these events have been reported voluntarily during clinical practice, estimates of frequency cannot be made and causal relationships between protease inhibitor therapy and these events have not been established.

5.6 Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in patients treated with combination antiretroviral therapy, including LEXIVA. During the initial phase of combination antiretroviral treatment, patients whose immune systems respond may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium* infection, cytomegalovirus, *Pneumocystis jirovecii* pneumonia [PCP], or tuberculosis), which may necessitate further evaluation and treatment.

5.7 Fat Redistribution

Redistribution/accumulation of body fat, including central obesity, dorsocervical fat enlargement (buffalo hump), peripheral wasting, facial wasting, breast enlargement, and “cushingoid appearance,” have been observed in patients receiving antiretroviral therapy, including LEXIVA. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been established.

5.8 Lipid Elevations

Treatment with LEXIVA plus ritonavir has resulted in increases in the concentration of triglycerides and cholesterol [*see Adverse Reactions (6)*]. Triglyceride and cholesterol testing should be performed prior to initiating therapy with LEXIVA and at periodic intervals during therapy. Lipid disorders should be managed as clinically appropriate [*see Drug Interactions (7)*].

5.9 Hemolytic Anemia

Acute hemolytic anemia has been reported in a patient treated with amprenavir.

5.10 Patients With Hemophilia

There have been reports of spontaneous bleeding in patients with hemophilia A and B treated with protease inhibitors. In some patients, additional factor VIII was required. In many of the reported cases, treatment with protease inhibitors was continued or restarted. A causal relationship between protease inhibitor therapy and these episodes has not been established.

5.11 Nephrolithiasis

Cases of nephrolithiasis were reported during postmarketing surveillance in HIV-infected patients receiving LEXIVA. Because these events were reported voluntarily during clinical practice, estimates of frequency cannot be made. If signs or symptoms of nephrolithiasis occur, temporary interruption or discontinuation of therapy may be considered.

5.12 Resistance/Cross-Resistance

Because the potential for HIV cross-resistance among protease inhibitors has not been fully explored, it is unknown what effect therapy with LEXIVA will have on the activity of subsequently administered protease inhibitors. LEXIVA has been studied in patients who have experienced treatment failure with protease inhibitors [*see Clinical Studies (14.2)*].

6 ADVERSE REACTIONS

- Severe or life-threatening skin reactions have been reported with the use of LEXIVA [*see Warnings and Precautions (5.2)*].

- The most common moderate to severe adverse reactions in clinical studies of LEXIVA were diarrhea, rash, nausea, vomiting, and headache.
- Treatment discontinuation due to adverse events occurred in 6.4% of patients receiving LEXIVA and in 5.9% of patients receiving comparator treatments. The most common adverse reactions leading to discontinuation of LEXIVA (incidence less than or equal to 1% of patients) included diarrhea, nausea, vomiting, AST increased, ALT increased, and rash.

6.1 Clinical Trials

Adults: The data for the 3 active-controlled clinical trials described below reflect exposure of 700 HIV-1 infected patients to LEXIVA Tablets, including 599 patients exposed to LEXIVA for greater than 24 weeks, and 409 patients exposed for greater than 48 weeks. The population age ranged from 17 to 72 years. Of these patients, 26% were female, 51% Caucasian, 31% black, 16% American Hispanic, and 70% were antiretroviral-naive. Sixty-one percent received LEXIVA 1,400 mg once daily plus ritonavir 200 mg once daily, 24% received LEXIVA 1,400 mg twice daily, and 15% received LEXIVA 700 mg twice daily plus ritonavir 100 mg twice daily.

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Selected adverse reactions reported during the clinical efficacy studies of LEXIVA are shown in Tables 2 and 3. Each table presents adverse reactions of moderate or severe intensity in patients treated with combination therapy for up to 48 weeks.

Table 2. Selected Moderate/Severe Clinical Adverse Reactions Reported in Greater Than or Equal to 2% of Antiretroviral-Naive Adult Patients

Adverse Reaction	APV30001 ^a		APV30002 ^a	
	LEXIVA 1,400 mg b.i.d. (n = 166)	Nelfinavir 1,250 mg b.i.d. (n = 83)	LEXIVA 1,400 mg q.d./ Ritonavir 200 mg q.d. (n = 322)	Nelfinavir 1,250 mg b.i.d. (n = 327)
Gastrointestinal				
Diarrhea	5%	18%	10%	18%
Nausea	7%	4%	7%	5%
Vomiting	2%	4%	6%	4%
Abdominal pain	1%	0%	2%	2%
Skin				
Rash	8%	2%	3%	2%
General disorders				
Fatigue	2%	1%	4%	2%
Nervous system				
Headache	2%	4%	3%	3%

^aAll patients also received abacavir and lamivudine twice daily.

Table 3. Selected Moderate/Severe Clinical Adverse Reactions Reported in Greater Than or Equal to 2% of Protease Inhibitor-Experienced Adult Patients (Study APV30003)

Adverse Reaction	LEXIVA 700 mg b.i.d./ Ritonavir 100 mg b.i.d. ^a (n = 106)	Lopinavir 400 mg b.i.d./ Ritonavir 100 mg b.i.d. ^a (n = 103)
Gastrointestinal		
Diarrhea	13%	11%
Nausea	3%	9%
Vomiting	3%	5%
Abdominal pain	<1%	2%
Skin		
Rash	3%	0%
Nervous system		
Headache	4%	2%

^aAll patients also received 2 reverse transcriptase inhibitors.

Skin rash (without regard to causality) occurred in approximately 19% of patients treated with LEXIVA in the pivotal efficacy studies. Rashes were usually maculopapular and of mild or moderate intensity, some with pruritus. Rash had a median onset of 11 days after initiation of

LEXIVA and had a median duration of 13 days. Skin rash led to discontinuation of LEXIVA in less than 1% of patients. In some patients with mild or moderate rash, dosing with LEXIVA was often continued without interruption; if interrupted, reintroduction of LEXIVA generally did not result in rash recurrence.

The percentages of patients with Grade 3 or 4 laboratory abnormalities in the clinical efficacy studies of LEXIVA are presented in Tables 4 and 5.

Table 4. Grade 3/4 Laboratory Abnormalities Reported in Greater Than or Equal to 2% of Antiretroviral-Naive Adult Patients in Studies APV30001 and APV30002

Laboratory Abnormality	APV30001 ^a		APV30002 ^a	
	LEXIVA 1,400 mg b.i.d. (n = 166)	Nelfinavir 1,250 mg b.i.d. (n = 83)	LEXIVA 1,400 mg q.d./ Ritonavir 200 mg q.d. (n = 322)	Nelfinavir 1,250 mg b.i.d. (n = 327)
ALT (>5 x ULN)	6%	5%	8%	8%
AST (>5 x ULN)	6%	6%	6%	7%
Serum lipase (>2 x ULN)	8%	4%	6%	4%
Triglycerides ^b (>750 mg/dL)	0%	1%	6%	2%
Neutrophil count, absolute (<750 cells/mm ³)	3%	6%	3%	4%

^aAll patients also received abacavir and lamivudine twice daily.

^bFasting specimens.

ULN = Upper limit of normal.

The incidence of Grade 3 or 4 hyperglycemia in antiretroviral-naive patients who received LEXIVA in the pivotal studies was less than 1%.

Table 5. Grade 3/4 Laboratory Abnormalities Reported in Greater Than or Equal to 2% of Protease Inhibitor-Experienced Adult Patients in Study APV30003

Laboratory Abnormality	LEXIVA 700 mg b.i.d./ Ritonavir 100 mg b.i.d. ^a (n = 104)	Lopinavir 400 mg b.i.d./ Ritonavir 100 mg b.i.d. ^a (n = 103)
Triglycerides ^b (>750 mg/dL)	11% ^c	6% ^c
Serum lipase (>2 x ULN)	5%	12%
ALT (>5 x ULN)	4%	4%
AST (>5 x ULN)	4%	2%
Glucose (>251 mg/dL)	2% ^c	2% ^c

^aAll patients also received 2 reverse transcriptase inhibitors.

^bFasting specimens.

^cn = 100 for LEXIVA plus ritonavir, n = 98 for lopinavir plus ritonavir.

ULN = Upper limit of normal.

Pediatric Patients: LEXIVA with and without ritonavir was studied in 144 pediatric patients aged 2 to 18 years in 2 open-label studies. Safety information from 75 pediatric patients receiving LEXIVA twice daily with or without ritonavir follows.

All adverse events regardless of causality, all drug-related adverse events, and all laboratory events occurred with similar frequency in pediatrics compared with adults, with the exception of vomiting. Vomiting, regardless of causality, occurred more frequently among pediatric patients receiving LEXIVA twice daily with ritonavir ([30%] all aged between 2 and 18 years) and without ritonavir ([56%] all aged between 2 and 5 years) compared with adults receiving LEXIVA twice daily with ritonavir (10%) and without ritonavir (16%). The median duration of drug-related vomiting episodes was 1 day (range: 1 to 62 days). Vomiting required temporary dose interruptions in 4 pediatric patients and was treatment-limiting in 1 pediatric patient, all of whom were receiving LEXIVA twice daily with ritonavir.

6.2 Postmarketing Experience

In addition to adverse reactions reported from clinical trials, the following reactions have been identified during post-approval use of LEXIVA. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. These reactions have been chosen for inclusion due to a combination of their seriousness, frequency of reporting, or potential causal connection to LEXIVA.

Cardiac Disorders: Myocardial infarction.

Metabolism and Nutrition Disorders: Hypercholesterolemia.

Nervous System Disorders: Oral paresthesia.

Skin and Subcutaneous Tissue Disorders: Angioedema.

Urogenital: Nephrolithiasis.

7 DRUG INTERACTIONS

See also Contraindications (4), Clinical Pharmacology (12.3).

If LEXIVA is used in combination with ritonavir, see full prescribing information for ritonavir for additional information on drug interactions.

7.1 CYP Inhibitors and Inducers

Amprenavir, the active metabolite of fosamprenavir, is an inhibitor of cytochrome P450 3A4 metabolism and therefore should not be administered concurrently with medications with narrow therapeutic windows that are substrates of CYP3A4. Data also suggest that amprenavir induces CYP3A4.

Amprenavir is metabolized by CYP3A4. Coadministration of LEXIVA and drugs that induce CYP3A4, such as rifampin, may decrease amprenavir concentrations and reduce its therapeutic effect. Coadministration of LEXIVA and drugs that inhibit CYP3A4 may increase amprenavir concentrations and increase the incidence of adverse effects.

The potential for drug interactions with LEXIVA changes when LEXIVA is coadministered with the potent CYP3A4 inhibitor ritonavir. The magnitude of CYP3A4-mediated drug interactions (effect on amprenavir or effect on coadministered drug) may change when LEXIVA is coadministered with ritonavir. Because ritonavir is a CYP2D6 inhibitor, clinically significant interactions with drugs metabolized by CYP2D6 are possible when coadministered with LEXIVA plus ritonavir.

There are other agents that may result in serious and/or life-threatening drug interactions [*see Contraindications (4)*].

7.2 Drugs That Should Not Be Coadministered With LEXIVA

See Contraindications (4).

7.3 Established and Other Potentially Significant Drug Interactions

Table 6 provides a listing of established or potentially clinically significant drug interactions. Information in the table applies to LEXIVA with or without ritonavir, unless otherwise indicated.

Table 6. Established and Other Potentially Significant Drug Interactions

Concomitant Drug Class: Drug Name	Effect on Concentration of Amprenavir or Concomitant Drug	Clinical Comment
<i>HIV-Antiviral Agents</i>		
Non-nucleoside reverse transcriptase inhibitor: Efavirenz ^a	LEXIVA: ↓Amprenavir LEXIVA/ritonavir: ↓Amprenavir	Appropriate doses of the combinations with respect to safety and efficacy have not been established. An additional 100 mg/day (300 mg total) of ritonavir is recommended when efavirenz is administered with LEXIVA/ritonavir once daily. No change in the ritonavir dose is required when efavirenz is administered with LEXIVA plus ritonavir twice daily.
Non-nucleoside reverse transcriptase inhibitor: Nevirapine ^a	LEXIVA: ↓Amprenavir ↑Nevirapine LEXIVA/ritonavir: ↓Amprenavir ↑Nevirapine	Coadministration of nevirapine and LEXIVA without ritonavir is not recommended. No dosage adjustment required when nevirapine is administered with LEXIVA/ritonavir twice daily. The combination of nevirapine administered with LEXIVA/ritonavir once-daily regimen has not been studied.
HIV protease inhibitor: Atazanavir ^a	LEXIVA: Interaction has not been evaluated. LEXIVA/ritonavir: ↓Atazanavir ↔Amprenavir	Appropriate doses of the combinations with respect to safety and efficacy have not been established.

HIV protease inhibitors: Indinavir ^a , nelfinavir ^a	LEXIVA: ↑Amprenavir Effect on indinavir and nelfinavir is not well established. LEXIVA/ritonavir: Interaction has not been evaluated.	Appropriate doses of the combinations with respect to safety and efficacy have not been established.
HIV protease inhibitors: Lopinavir/ritonavir ^a	↓Amprenavir ↓Lopinavir	An increased rate of adverse events has been observed. Appropriate doses of the combinations with respect to safety and efficacy have not been established.
HIV protease inhibitor: Saquinavir ^a	LEXIVA: ↓Amprenavir Effect on saquinavir is not well established. LEXIVA/ritonavir: Interaction has not been evaluated.	Appropriate doses of the combination with respect to safety and efficacy have not been established.
HIV integrase inhibitor: Raltegravir ^a	LEXIVA: ↓Amprenavir ↓Raltegravir LEXIVA/ritonavir: ↓Amprenavir ↓Raltegravir	Appropriate doses of the combination with respect to safety and efficacy have not been established [<i>see Clinical Pharmacology (12.3)</i>].
Other Agents		
Antiarrhythmics: Amiodarone, bepridil, lidocaine (systemic), and quinidine	↑Antiarrhythmics	Use with caution. Increased exposure may be associated with life-threatening reactions such as cardiac arrhythmias. Therapeutic concentration monitoring, if available, is recommended for antiarrhythmics.
Anticoagulant:		Concentrations of warfarin may be

Warfarin		affected. It is recommended that INR (international normalized ratio) be monitored.
Anticonvulsants: Carbamazepine, phenobarbital, phenytoin	LEXIVA: ↓Amprenavir	Use with caution. LEXIVA may be less effective due to decreased amprenavir plasma concentrations in patients taking these agents concomitantly.
Phenytoin ^a	LEXIVA/ritonavir: ↑Amprenavir ↓Phenytoin	Plasma phenytoin concentrations should be monitored and phenytoin dose should be increased as appropriate. No change in LEXIVA/ritonavir dose is recommended.
Antidepressant: Paroxetine, trazodone	↓Paroxetine ↑Trazodone	Coadministration of paroxetine with LEXIVA/ritonavir significantly decreased plasma levels of paroxetine. Any paroxetine dose adjustment should be guided by clinical effect (tolerability and efficacy). Concomitant use of trazodone and LEXIVA with or without ritonavir may increase plasma concentrations of trazodone. Adverse events of nausea, dizziness, hypotension, and syncope have been observed following coadministration of trazodone and ritonavir. If trazodone is used with a CYP3A4 inhibitor such as LEXIVA, the combination should be used with caution and a lower dose of trazodone should be considered.

<p>Antifungals: Ketoconazole^a, itraconazole</p>	<p>↑Ketoconazole ↑Itraconazole</p>	<p>Increase monitoring for adverse events. LEXIVA: Dose reduction of ketoconazole or itraconazole may be needed for patients receiving more than 400 mg ketoconazole or itraconazole per day. LEXIVA/ritonavir: High doses of ketoconazole or itraconazole (greater than 200 mg/day) are not recommended.</p>
<p>Anti-gout: Colchicine</p>	<p>↑Colchicine</p>	<p>Patients with renal or hepatic impairment should not be given colchicine with LEXIVA/ritonavir.</p> <p>LEXIVA/ritonavir and coadministration of colchicine:</p> <p>Treatment of gout flares: 0.6 mg (1 tablet) x 1 dose, followed by 0.3 mg (half tablet) 1 hour later. Dose to be repeated no earlier than 3 days.</p> <p>Prophylaxis of gout flares: If the original regimen was 0.6 mg twice a day, the regimen should be adjusted to 0.3 mg once a day. If the original regimen was 0.6 mg once a day, the regimen should be adjusted to 0.3 mg once every other day.</p> <p>Treatment of familial Mediterranean fever (FMF): Maximum daily dose of 0.6 mg (may be given as 0.3 mg twice a day).</p> <p>LEXIVA and coadministration of colchicine:</p> <p>Treatment of gout flares: 1.2 mg (2 tablets) x 1 dose. Dose to be</p>

		<p>repeated no earlier than 3 days.</p> <p>Prophylaxis of gout flares: If the original regimen was 0.6 mg twice a day, the regimen should be adjusted to 0.3 mg twice a day or 0.6 mg once a day. If the original regimen was 0.6 mg once a day, the regimen should be adjusted to 0.3 mg once a day.</p> <p>Treatment of FMF: Maximum daily dose of 1.2 mg (may be given as 0.6 mg twice a day).</p>
<p>Antimycobacterial: Rifabutin^a</p>	<p>↑Rifabutin and rifabutin metabolite</p>	<p>A complete blood count should be performed weekly and as clinically indicated to monitor for neutropenia.</p> <p>LEXIVA: A dosage reduction of rifabutin by at least half the recommended dose is required.</p> <p>LEXIVA/ritonavir: Dosage reduction of rifabutin by at least 75% of the usual dose of 300 mg/day is recommended (a maximum dose of 150 mg every other day or 3 times per week).</p>
<p>Benzodiazepines: Alprazolam, clorazepate, diazepam, flurazepam</p>	<p>↑Benzodiazepines</p>	<p>Clinical significance is unknown. A decrease in benzodiazepine dose may be needed.</p>
<p>Calcium channel blockers: Diltiazem, felodipine, nifedipine, nicardipine, nimodipine, verapamil, amlodipine, nisoldipine, isradipine</p>	<p>↑Calcium channel blockers</p>	<p>Use with caution. Clinical monitoring of patients is recommended.</p>
<p>Corticosteroid: Dexamethasone</p>	<p>↓Amprenavir</p>	<p>Use with caution. LEXIVA may be less effective due to decreased amprenavir plasma concentrations.</p>
<p>Endothelin receptor</p>	<p>↑Bosentan</p>	<p>Coadministration of bosentan in patients</p>

<p>antagonists: Bosentan</p>		<p>on LEXIVA:</p> <p>In patients who have been receiving LEXIVA for at least 10 days, start bosentan at 62.5 mg once daily or every other day based upon individual tolerability.</p> <p>Coadministration of LEXIVA in patients on bosentan:</p> <p>Discontinue use of bosentan at least 36 hours prior to initiation of LEXIVA.</p> <p>After at least 10 days following the initiation of LEXIVA, resume bosentan at 62.5 mg once daily or every other day based upon individual tolerability.</p>
<p>Histamine H₂-receptor antagonists: Cimetidine, famotidine, nizatidine, ranitidine^a</p>	<p>LEXIVA: ↓Amprenavir</p> <p>LEXIVA/ritonavir: Interaction not evaluated</p>	<p>Use with caution. LEXIVA may be less effective due to decreased amprenavir plasma concentrations.</p>
<p>HMG-CoA reductase inhibitors: Atorvastatin^a, rosuvastatin</p>	<p>↑Atorvastatin ↑Rosuvastatin</p>	<p>Use the lowest possible dose of atorvastatin or rosuvastatin with careful monitoring, or consider other HMG-CoA reductase inhibitors such as fluvastatin or pravastatin.</p>
<p>Immunosuppressants: Cyclosporine, tacrolimus, rapamycin</p>	<p>↑Immunosuppressants</p>	<p>Therapeutic concentration monitoring is recommended for immunosuppressant agents.</p>
<p>Inhaled beta agonist: Salmeterol</p>	<p>↑Salmeterol</p>	<p>Concurrent administration of salmeterol with LEXIVA is not recommended. The combination may result in increased risk of cardiovascular adverse events associated with salmeterol, including QT prolongation, palpitations, and sinus tachycardia.</p>
<p>Inhaled/nasal steroid: Fluticasone</p>	<p>LEXIVA: ↑Fluticasone</p>	<p>Use with caution. Consider alternatives to fluticasone, particularly for long-term use.</p>

	LEXIVA/ritonavir: ↑Fluticasone	May result in significantly reduced serum cortisol concentrations. Systemic corticosteroid effects including Cushing's syndrome and adrenal suppression have been reported during postmarketing use in patients receiving ritonavir and inhaled or intranasally administered fluticasone. Coadministration of fluticasone and LEXIVA/ritonavir is not recommended unless the potential benefit to the patient outweighs the risk of systemic corticosteroid side effects.
Narcotic analgesic: Methadone	↓Methadone	Data suggest that the interaction is not clinically relevant; however, patients should be monitored for opiate withdrawal symptoms.
Oral contraceptives: Ethinyl estradiol/norethindrone ^a	LEXIVA: ↓Amprenavir ↓Ethinyl estradiol LEXIVA/ritonavir: ↓Ethinyl estradiol	Alternative methods of non-hormonal contraception are recommended. May lead to loss of virologic response. ^a Increased risk of transaminase elevations. No data are available on the use of LEXIVA/ritonavir with other hormonal therapies, such as hormone replacement therapy (HRT) for postmenopausal women.
PDE5 inhibitors: Sildenafil, tadalafil, vardenafil	↑Sildenafil ↑Tadalafil ↑Vardenafil	May result in an increase in PDE5 inhibitor-associated adverse events, including hypotension, syncope, visual disturbances, and priapism. <u>Use of PDE5 inhibitors for pulmonary arterial hypertension (PAH):</u> <ul style="list-style-type: none"> • Use of sildenafil (REVATIO) is contraindicated when used for the treatment of PAH-[see <i>Contraindications (4)</i>].

		<ul style="list-style-type: none"> • <u>The following dose adjustments are recommended for use of tadalafil (ADCIRCA™) with LEXIVA:</u> <p><u>Coadministration of ADCIRCA in patients on LEXIVA:</u></p> <p>In patients receiving LEXIVA for at least one week, start ADCIRCA at 20 mg once daily. Increase to 40 mg once daily based upon individual tolerability.</p> <p><u>Coadministration of LEXIVA in patients on ADCIRCA:</u></p> <p>Avoid use of ADCIRCA during the initiation of LEXIVA. Stop ADCIRCA at least 24 hours prior to starting LEXIVA. After at least one week following the initiation of LEXIVA, resume ADCIRCA at 20 mg once daily. Increase to 40 mg once daily based upon individual tolerability.</p> <p><u>Use of PDE5 inhibitors for erectile dysfunction:</u></p> <p>LEXIVA: Sildenafil: 25 mg every 48 hours. Tadalafil: no more than 10 mg every 72 hours. Vardenafil: no more than 2.5 mg every 24 hours.</p> <p>LEXIVA/ritonavir: Sildenafil: 25 mg every 48 hours. Tadalafil: no more than 10 mg every 72 hours. Vardenafil: no more than 2.5 mg every 72 hours.</p>
--	--	---

		Use with increased monitoring for adverse events.
Proton pump inhibitors: Esomeprazole ^a , lansoprazole, omeprazole, pantoprazole, rabeprazole	LEXIVA: ↔Amprenavir ↑Esomeprazole LEXIVA/ritonavir: ↔Amprenavir ↔Esomeprazole	Proton pump inhibitors can be administered at the same time as a dose of LEXIVA with no change in plasma amprenavir concentrations.
Tricyclic antidepressants: Amitriptyline, imipramine	↑Tricyclics	Therapeutic concentration monitoring is recommended for tricyclic antidepressants.

^a See *Clinical Pharmacology (12.3) Tables 10, 11, 12, or 13 for magnitude of interaction.*

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C. Embryo/fetal development studies were conducted in rats (dosed from day 6 to day 17 of gestation) and rabbits (dosed from day 7 to day 20 of gestation). Administration of fosamprenavir to pregnant rats and rabbits produced no major effects on embryo-fetal development; however, the incidence of abortion was increased in rabbits that were administered fosamprenavir. Systemic exposures ($AUC_{0-24 \text{ hr}}$) to amprenavir at these dosages were 0.8 (rabbits) to 2 (rats) times the exposures in humans following administration of the maximum recommended human dose (MRHD) of fosamprenavir alone or 0.3 (rabbits) to 0.7 (rats) times the exposures in humans following administration of the MRHD of fosamprenavir in combination with ritonavir. In contrast, administration of amprenavir was associated with abortions and an increased incidence of minor skeletal variations resulting from deficient ossification of the femur, humerus, and trochlea, in pregnant rabbits at the tested dose; approximately one-twentieth the exposure seen at the recommended human dose.

The mating and fertility of the F₁ generation born to female rats given fosamprenavir was not different from control animals; however, fosamprenavir did cause a reduction in both pup survival and body weights. Surviving F₁ female rats showed an increased time to successful mating, an increased length of gestation, a reduced number of uterine implantation sites per litter, and reduced gestational body weights compared with control animals. Systemic exposure ($AUC_{0-24 \text{ hr}}$) to amprenavir in the F₀ pregnant rats was approximately 2 times higher than exposures in humans following administration of the MRHD of fosamprenavir alone or approximately the same as those seen in humans following administration of the MRHD of fosamprenavir in combination with ritonavir.

There are no adequate and well-controlled studies in pregnant women. LEXIVA should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Antiretroviral Pregnancy Registry: To monitor maternal-fetal outcomes of pregnant women exposed to LEXIVA, an Antiretroviral Pregnancy Registry has been established. Physicians are encouraged to register patients by calling 1-800-258-4263.

8.3 Nursing Mothers

The Centers for Disease Control and Prevention recommend that HIV-infected mothers not breastfeed their infants to avoid risking postnatal transmission of HIV. Although it is not known if amprenavir is excreted in human milk, amprenavir is secreted into the milk of lactating rats. Because of both the potential for HIV transmission and the potential for serious adverse reactions in nursing infants, mothers should be instructed not to breastfeed if they are receiving LEXIVA.

8.4 Pediatric Use

The safety, pharmacokinetic profile, and virologic response of LEXIVA Oral Suspension and Tablets were evaluated in pediatric patients aged 2 to 18 years in 2 open-label studies [*see Clinical Studies (14.3)*]. No data are available for pediatric patients younger than 2 years.

The adverse reaction profile seen in pediatrics was similar to that seen in adults. Vomiting, regardless of causality, was more frequent in pediatrics than in adults [*see Adverse Reactions (6.1)*].

8.5 Geriatric Use

Clinical studies of LEXIVA did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger adults. In general, dose selection for an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

8.6 Hepatic Impairment

Amprenavir is principally metabolized by the liver; therefore, caution should be exercised when administering LEXIVA to patients with hepatic impairment because amprenavir concentrations may be increased [*see Clinical Pharmacology (12.3)*]. Patients with impaired hepatic function receiving LEXIVA with or without concurrent ritonavir require dose reduction [*see Dosage and Administration (2.3)*].


10 OVERDOSAGE

In a healthy volunteer repeat-dose pharmacokinetic study evaluating high-dose combinations of LEXIVA plus ritonavir, an increased frequency of Grade 2/3 ALT elevations (greater than 2.5 x ULN) was observed with LEXIVA 1,400 mg twice daily plus ritonavir 200 mg twice daily (4 of 25 subjects). Concurrent Grade 1/2 elevations in AST (greater than 1.25 x ULN) were noted in 3 of these 4 subjects. These transaminase elevations resolved following discontinuation of dosing.

There is no known antidote for LEXIVA. It is not known whether amprenavir can be removed by peritoneal dialysis or hemodialysis. If overdose occurs, the patient should be monitored for evidence of toxicity and standard supportive treatment applied as necessary.

11 DESCRIPTION

LEXIVA (fosamprenavir calcium) is a prodrug of amprenavir, an inhibitor of HIV protease. The chemical name of fosamprenavir calcium is (3*S*)-tetrahydrofuran-3-yl (1*S*,2*R*)-3-[[[4-aminophenyl) sulfonyl](isobutyl)amino]-1-benzyl-2-(phosphonoxy) propylcarbamate monocalcium salt. Fosamprenavir calcium is a single stereoisomer with the (3*S*)(1*S*,2*R*) configuration. It has a molecular formula of C₂₅H₃₄CaN₃O₉PS and a molecular weight of 623.7. It has the following structural formula:



Fosamprenavir calcium is a white to cream-colored solid with a solubility of approximately 0.31 mg/mL in water at 25°C.

LEXIVA Tablets are available for oral administration in a strength of 700 mg of fosamprenavir as fosamprenavir calcium (equivalent to approximately 600 mg of amprenavir). Each 700-mg tablet contains the inactive ingredients colloidal silicon dioxide, croscarmellose sodium, magnesium stearate, microcrystalline cellulose, and povidone K30. The tablet film-coating contains the inactive ingredients hypromellose, iron oxide red, titanium dioxide, and triacetin.

LEXIVA Oral Suspension is available in a strength of 50 mg/mL of fosamprenavir as fosamprenavir calcium equivalent to approximately 43 mg of amprenavir. LEXIVA Oral Suspension is a white to off-white suspension with a grape-bubblegum-peppermint flavor. Each one milliliter (1 mL) contains the inactive ingredients artificial grape-bubblegum flavor, calcium chloride dihydrate, hypromellose, methylparaben, natural peppermint flavor, polysorbate 80, propylene glycol, propylparaben, purified water, and sucralose.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Fosamprenavir is an antiviral agent [see *Clinical Pharmacology* (12.4)].

12.3 Pharmacokinetics

The pharmacokinetic properties of amprenavir after administration of LEXIVA, with or without ritonavir, have been evaluated in both healthy adult volunteers and in HIV-infected

patients; no substantial differences in steady-state amprenavir concentrations were observed between the 2 populations.

The pharmacokinetic parameters of amprenavir after administration of LEXIVA (with and without concomitant ritonavir) are shown in Table 7.


Table 7. Geometric Mean (95% CI) Steady-State Plasma Amprenavir Pharmacokinetic Parameters in Adults

Regimen	C _{max} (mcg/mL)	T _{max} (hours) ^a	AUC ₂₄ (mcg•hr/mL)	C _{min} (mcg/mL)
LEXIVA 1,400 mg b.i.d.	4.82 (4.06-5.72)	1.3 (0.8-4.0)	33.0 (27.6-39.2)	0.35 (0.27-0.46)
LEXIVA 1,400 mg q.d. plus Ritonavir 200 mg q.d.	7.24 (6.32-8.28)	2.1 (0.8-5.0)	69.4 (59.7-80.8)	1.45 (1.16-1.81)
LEXIVA 1,400 mg q.d. plus Ritonavir 100 mg q.d.	7.93 (7.25-8.68)	1.5 (0.75-5.0)	66.4 (61.1-72.1)	0.86 (0.74-1.01)
LEXIVA 700 mg b.i.d. plus Ritonavir 100 mg b.i.d.	6.08 (5.38-6.86)	1.5 (0.75-5.0)	79.2 (69.0-90.6)	2.12 (1.77-2.54)

^aData shown are median (range).

The mean plasma amprenavir concentrations of the dosing regimens over the dosing intervals are displayed in Figure 1.

Figure 1. Mean (\pm SD) Steady-State Plasma Amprenavir Concentrations and Mean IC_{50} Values Against HIV from Protease Inhibitor-Naive Patients (in the Absence of Human Serum)



Absorption and Bioavailability: After administration of a single dose of LEXIVA to HIV-1-infected patients, the time to peak amprenavir concentration (T_{max}) occurred between 1.5 and 4 hours (median 2.5 hours). The absolute oral bioavailability of amprenavir after administration of LEXIVA in humans has not been established.

After administration of a single 1,400-mg dose in the fasted state, LEXIVA Oral Suspension (50 mg/mL) and LEXIVA Tablets (700 mg) provided similar amprenavir exposures (AUC), however, the C_{max} of amprenavir after administration of the suspension formulation was 14.5% higher compared with the tablet.

Effects of Food on Oral Absorption: Administration of a single 1,400-mg dose of LEXIVA Tablets in the fed state (standardized high-fat meal: 967 kcal, 67 grams fat, 33 grams protein, 58 grams carbohydrate) compared with the fasted state was associated with no significant changes in amprenavir C_{max} , T_{max} , or $AUC_{0-\infty}$ [see *Dosage and Administration (2)*].

Administration of a single 1,400-mg dose of LEXIVA Oral Suspension in the fed state (standardized high-fat meal: 967 kcal, 67 grams fat, 33 grams protein, 58 grams carbohydrate) compared with the fasted state was associated with a 46% reduction in C_{max} , a 0.72-hour delay in T_{max} , and a 28% reduction in amprenavir $AUC_{0-\infty}$.

Distribution: In vitro, amprenavir is approximately 90% bound to plasma proteins, primarily to α_1 -acid glycoprotein. In vitro, concentration-dependent binding was observed

over the concentration range of 1 to 10 mcg/mL, with decreased binding at higher concentrations. The partitioning of amprenavir into erythrocytes is low, but increases as amprenavir concentrations increase, reflecting the higher amount of unbound drug at higher concentrations.

Metabolism: After oral administration, fosamprenavir is rapidly and almost completely hydrolyzed to amprenavir and inorganic phosphate prior to reaching the systemic circulation. This occurs in the gut epithelium during absorption. Amprenavir is metabolized in the liver by the cytochrome P450 3A4 (CYP3A4) enzyme system. The 2 major metabolites result from oxidation of the tetrahydrofuran and aniline moieties. Glucuronide conjugates of oxidized metabolites have been identified as minor metabolites in urine and feces.

Amprenavir is both a substrate for and inducer of P-glycoprotein.

Elimination: Excretion of unchanged amprenavir in urine and feces is minimal. Unchanged amprenavir in urine accounts for approximately 1% of the dose; unchanged amprenavir was not detectable in feces. Approximately 14% and 75% of an administered single dose of ¹⁴C-amprenavir can be accounted for as metabolites in urine and feces, respectively. Two metabolites accounted for greater than 90% of the radiocarbon in fecal samples. The plasma elimination half-life of amprenavir is approximately 7.7 hours.

Special Populations: Hepatic Impairment: The pharmacokinetics of amprenavir have been studied after the administration of LEXIVA in combination with ritonavir to adult HIV-1-infected patients with mild, moderate, and severe hepatic impairment. Following 2 weeks of dosing with LEXIVA plus ritonavir, the AUC of amprenavir was increased by approximately 22% in patients with mild hepatic impairment, by approximately 70% in patients with moderate hepatic impairment, and by approximately 80% in patients with severe hepatic impairment compared with HIV-1-infected patients with normal hepatic function. Protein binding of amprenavir was decreased in patients with hepatic impairment. The unbound fraction at 2 hours (approximate C_{max}) ranged between a decrease of -7% to an increase of 57% while the unbound fraction at the end of the dosing interval (C_{min}) increased from 50% to 102% [*see Dosage and Administration (2.3)*].

The pharmacokinetics of amprenavir have been studied after administration of amprenavir given as AGENERASE[®] Capsules to adult patients with hepatic impairment. Following administration of a single 600-mg oral dose, the AUC of amprenavir was increased by approximately 2.5-fold in patients with moderate cirrhosis and by approximately 4.5-fold in patients with severe cirrhosis compared with healthy volunteers [*see Dosage and Administration (2.3)*].

Renal Impairment: The impact of renal impairment on amprenavir elimination in adult patients has not been studied. The renal elimination of unchanged amprenavir represents approximately 1% of the administered dose; therefore, renal impairment is not expected to significantly impact the elimination of amprenavir.

Pediatric Patients: The pharmacokinetics of amprenavir after administration of LEXIVA Oral Suspension and LEXIVA Tablets, with or without ritonavir, have been evaluated

in 124 patients aged 2 to 18 years. Pharmacokinetic parameters for LEXIVA administered with food and with or without ritonavir in this patient population are provided in Tables 8 and 9 below.

Table 8. Geometric Mean (95% CI) Steady-State Plasma Amprenavir Pharmacokinetic Parameters in Pediatric Patients Receiving LEXIVA 30 mg/kg Twice Daily

Parameter	2 to 5 Years	
	n	LEXIVA 30 mg/kg b.i.d.
AUC ₍₂₄₎ (mcg•hr/mL)	8	31.4 (13.7, 72.4)
C _{max} (mcg/mL)	8	5.00 (1.95, 12.8)
C _{min} (mcg/mL)	17	0.454 (0.342, 0.604)

Table 9. Geometric Mean (95% CI) Steady-State Plasma Amprenavir Pharmacokinetic Parameters in Pediatric and Adolescent Patients Receiving LEXIVA Plus Ritonavir Twice Daily

Parameter	6 to 11 Years		12 to 18 Years	
	n	LEXIVA 18 mg/kg plus Ritonavir 3 mg/kg b.i.d.	n	LEXIVA 700 mg plus Ritonavir 100 mg b.i.d.
AUC ₍₀₋₂₄₎ (mcg•hr/mL)	9	93.4 (67.8, 129)	8	58.8 (38.8, 89.0)
C _{max} (mcg/mL)	9	6.07 (4.40, 8.38)	8	4.33 (2.82, 6.65)
C _{min} (mcg/mL)	17	2.69 (2.15, 3.36)	24	1.61 (1.21, 2.15)

Geriatric Patients: The pharmacokinetics of amprenavir after administration of LEXIVA to patients older than 65 years have not been studied [see *Use in Specific Populations* (8.5)].

Gender: The pharmacokinetics of amprenavir after administration of LEXIVA do not differ between males and females.

Race: The pharmacokinetics of amprenavir after administration of LEXIVA do not differ between blacks and non-blacks.

Drug Interactions: [See *Contraindications* (4), *Warnings and Precautions* (5.1), *Drug Interactions* (7).]

Amprenavir, the active metabolite of fosamprenavir, is metabolized in the liver by the cytochrome P450 enzyme system. Amprenavir inhibits CYP3A4. Data also suggest that amprenavir induces CYP3A4. Caution should be used when coadministering medications that

are substrates, inhibitors, or inducers of CYP3A4, or potentially toxic medications that are metabolized by CYP3A4. Amprenavir does not inhibit CYP2D6, CYP1A2, CYP2C9, CYP2C19, CYP2E1, or uridine glucuronosyltransferase (UDPGT).

Drug interaction studies were performed with LEXIVA and other drugs likely to be coadministered or drugs commonly used as probes for pharmacokinetic interactions. The effects of coadministration on AUC, C_{\max} , and C_{\min} values are summarized in Table 10 (effect of other drugs on amprenavir) and Table 12 (effect of LEXIVA on other drugs). In addition, since LEXIVA delivers comparable amprenavir plasma concentrations as AGENERASE, drug interaction data derived from studies with AGENERASE are provided in Tables 11 and 13. For information regarding clinical recommendations, *see Drug Interactions (7)*.

Table 10. Drug Interactions: Pharmacokinetic Parameters for Amprenavir After Administration of LEXIVA in the Presence of the Coadministered Drug(s)

Coadministered Drug(s) and Dose(s)	Dose of LEXIVA ^a	n	% Change in Amprenavir Pharmacokinetic Parameters (90% CI)		
			C _{max}	AUC	C _{min}
Antacid (MAALOX TC [®]) 30 mL single dose	1,400 mg single dose	30	↓35 (↓24 to ↓42)	↓18 (↓9 to ↓26)	↑14 (↓7 to ↑39)
Atazanavir 300 mg q.d. for 10 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 10 days	22	↔	↔	↔
Atorvastatin 10 mg q.d. for 4 days	1,400 mg b.i.d. for 2 weeks	16	↓18 (↓34 to ↑1)	↓27 (↓41 to ↓12)	↓12 (↓27 to ↓6)
Atorvastatin 10 mg q.d. for 4 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	16	↔	↔	↔
Efavirenz 600 mg q.d. for 2 weeks	1,400 mg q.d. plus ritonavir 200 mg q.d. for 2 weeks	16	↔	↓13 (↓30 to ↑7)	↓36 (↓8 to ↓56)
Efavirenz 600 mg q.d. plus additional ritonavir 100 mg q.d. for 2 weeks	1,400 mg q.d. plus ritonavir 200 mg q.d. for 2 weeks	16	↑18 (↑1 to ↑38)	↑11 (0 to ↑24)	↔
Efavirenz 600 mg q.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	16	↔	↔	↓17 (↓4 to ↓29)
Esomeprazole 20 mg q.d. for 2 weeks	1,400 mg b.i.d. for 2 weeks	25	↔	↔	↔
Esomeprazole 20 mg q.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	23	↔	↔	↔
Ethinyl estradiol/norethin- drone 0.035 mg/0.5 mg q.d. for 21 days	700 mg b.i.d. plus ritonavir ^b 100 mg b.i.d. for 21 days	25	↔ ^c	↔ ^c	↔ ^c

Ketoconazole ^d 200 mg q.d. for 4 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 4 days	15	↔	↔	↔
Lopinavir/ritonavir 533 mg/133 mg b.i.d.	1,400 mg b.i.d. for 2 weeks	18	↓13 ^e	↓26 ^e	↓42 ^e
Lopinavir/ritonavir 400 mg/100 mg b.i.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	18	↓58 (↓42 to ↓70)	↓63 (↓51 to ↓72)	↓65 (↓54 to ↓73)
Methadone 70 to 120 mg q.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	19	↔ ^c	↔ ^c	↔ ^c
Nevirapine 200 mg b.i.d. for 2 weeks ^f	1,400 mg b.i.d. for 2 weeks	17	↓25 (↓37 to ↓10)	↓33 (↓45 to ↓20)	↓35 (↓50 to ↓15)
Nevirapine 200 mg b.i.d. for 2 weeks ^f	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	17	↔	↓11 (↓23 to ↑3)	↓19 (↓32 to ↓4)
Phenytoin 300 mg q.d. for 10 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 10 days	13	↔	↑20 (↑8 to ↑34)	↑19 (↑6 to ↑33)
Raltegravir 400 mg b.i.d. for 14 days	1,400 mg b.i.d. for 14 days (fasted)	14	↓27 (↓46 to ↔)	↓36 (↓53 to ↓13)	↓43 ^g (↓59 to ↓21)
	1,400 mg b.i.d. for 14 days ^h	14	↓15 (↓27 to ↓1)	↓17 (↓27 to ↓6)	↓32 ^g (↓53 to ↓1)
	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 14 days (fasted)	14	↓14 (↓39 to ↑20)	↓17 (↓38 to ↑12)	↓20 ^g (↓45 to ↑17)
	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 14 days ^h	12	↓25 (↓42 to ↓2)	↓25 (↓44 to ↔)	↓33 ^g (↓52 to ↓7)
Raltegravir 400 mg b.i.d. for 14 days	1,400 mg q.d. plus ritonavir 100 mg q.d. for 14 days (fasted)	13	↓18 (↓34 to ↔)	↓24 (↓41 to ↔)	↓50 ^g (↓64 to ↓31)

	1,400 mg q.d. plus ritonavir 100 mg q.d for 14 days ^h	14	↑27 (↓1 to ↑62)	↑13 (↓7 to ↑38)	↓17 ^g (↓45 to ↑26)
Ranitidine 300 mg single dose (administered 1 hour before fosamprenavir)	1,400 mg single dose	30	↓51 (↓43 to ↓58)	↓30 (↓22 to ↓37)	↔ (↓19 to ↑21)
Rifabutin 150 mg q.o.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	15	↑36 ^c (↑18 to ↑55)	↑35 ^c (↑17 to ↑56)	↑17 ^c (↓1 to ↑39)
Tenofovir 300 mg q.d. for 4 to 48 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 4 to 48 weeks	45	NA	NA	↔ ⁱ
Tenofovir 300 mg q.d. for 4 to 48 weeks	1,400 mg q.d. plus ritonavir 200 mg q.d. for 4 to 48 weeks	60	NA	NA	↔ ⁱ

^a Concomitant medication is also shown in this column where appropriate.

^b Ritonavir C_{max}, AUC, and C_{min} increased by 63%, 45%, and 13%, respectively, compared with historical control.

^c Compared with historical control.

^d Patients were receiving LEXIVA/ritonavir for 10 days prior to the 4-day treatment period with both ketoconazole and LEXIVA/ritonavir.

^e Compared with LEXIVA 700 mg/ritonavir 100 mg b.i.d. for 2 weeks.

^f Patients were receiving nevirapine for at least 12 weeks prior to study.

^g C_{last} (C_{12 hr} or C_{24 hr}).

^h Doses of LEXIVA and raltegravir were given with food on PK sampling days and without regard to food all other days.

ⁱ Compared with parallel control group.

↑= Increase; ↓= Decrease; ↔ = No change (↑ or ↓ less than or equal to 10%), NA = Not applicable.

Table 11. Drug Interactions: Pharmacokinetic Parameters for Amprenavir After Administration of AGENERASE in the Presence of the Coadministered Drug(s)

Coadministered Drug(s) and Dose(s)	Dose of AGENERASE ^a	n	% Change in Amprenavir Pharmacokinetic Parameters (90% CI)		
			C _{max}	AUC	C _{min}
Abacavir 300 mg b.i.d. for 2 to 3 weeks	900 mg b.i.d. for 2 to 3 weeks	4	↔ ^a	↔ ^a	↔ ^a
Clarithromycin 500 mg b.i.d. for 4 days	1,200 mg b.i.d. for 4 days	12	↑15 (↑1 to ↑31)	↑18 (↑8 to ↑29)	↑39 (↑31 to ↑47)
Delavirdine 600 mg b.i.d. for 10 days	600 mg b.i.d. for 10 days	9	↑40 ^b	↑130 ^b	↑125 ^b
Ethinyl estradiol/norethindrone 0.035 mg/1 mg for 1 cycle	1,200 mg b.i.d. for 28 days	10	↔	↓22 (↓35 to ↓8)	↓20 (↓41 to ↑8)
Indinavir 800 mg t.i.d. for 2 weeks (fasted)	750 or 800 mg t.i.d. for 2 weeks (fasted)	9	↑18 (↑13 to ↑58)	↑33 (↑2 to ↑73)	↑25 (↓27 to ↑116)
Ketoconazole 400 mg single dose	1,200 mg single dose	12	↓16 (↓25 to ↓6)	↑31 (↑20 to ↑42)	NA
Lamivudine 150 mg single dose	600 mg single dose	11	↔	↔	NA
Methadone 44 to 100 mg q.d. for >30 days	1,200 mg b.i.d. for 10 days	16	↓27 ^c	↓30 ^c	↓25 ^c
Nelfinavir 750 mg t.i.d. for 2 weeks (fed)	750 or 800 mg t.i.d. for 2 weeks (fed)	6	↓14 (↓38 to ↑20)	↔	↑189 (↑52 to ↑448)
Rifabutin 300 mg q.d. for 10 days	1,200 mg b.i.d. for 10 days	5	↔	↓15 (↓28 to 0)	↓15 (↓38 to ↑17)
Rifampin 300 mg q.d. for 4 days	1,200 mg b.i.d. for 4 days	11	↓70 (↓76 to ↓62)	↓82 (↓84 to ↓78)	↓92 (↓95 to ↓89)
Saquinavir 800 mg t.i.d. for 2 weeks (fed)	750 or 800 mg t.i.d. for 2 weeks (fed)	7	↓37 (↓54 to ↓14)	↓32 (↓49 to ↓9)	↓14 (↓52 to ↑54)
Zidovudine 300 mg single dose	600 mg single dose	12	↔	↑13 (↓2 to ↑31)	NA

^a Compared with parallel control group.

^b Median percent change; confidence interval not reported.

^c Compared with historical data.

↑ = Increase; ↓ = Decrease; ↔ = No change (↑ or ↓ less than 10%); NA = C_{min} not calculated for single-dose study.

Table 12. Drug Interactions: Pharmacokinetic Parameters for Coadministered Drug in the Presence of Amprenavir After Administration of LEXIVA

Coadministered Drug(s) and Dose(s)	Dose of LEXIVA ^a	n	% Change in Pharmacokinetic Parameters of Coadministered Drug (90% CI)		
			C _{max}	AUC	C _{min}
Atazanavir 300 mg q.d. for 10 days ^b	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 10 days	21	↓24 (↓39 to ↓6)	↓22 (↓34 to ↓9)	↔
Atorvastatin 10 mg q.d. for 4 days	1,400 mg b.i.d. for 2 weeks	16	↑304 (↑205 to ↑437)	↑130 (↑100 to ↑164)	↓10 (↓27 to ↑12)
Atorvastatin 10 mg q.d. for 4 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	16	↑184 (↑126 to ↑257)	↑153 (↑115 to ↑199)	↑73 (↑45 to ↑108)
Esomeprazole 20 mg q.d. for 2 weeks	1,400 mg b.i.d. for 2 weeks	25	↔	↑55 (↑39 to ↑73)	ND
Esomeprazole 20 mg q.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	23	↔	↔	ND
Ethinyl estradiol ^c 0.035 mg q.d. for 21 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 21 days	25	↓28 (↓21 to ↓35)	↓37 (↓30 to ↓42)	ND
Ketoconazole ^d 200 mg q.d. for 4 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 4 days	15	↑25 (↑0 to ↑56)	↑169 (↑108 to ↑248)	ND
Lopinavir/ritonavir ^e 533 mg/133 mg b.i.d. for 2 weeks	1,400 mg b.i.d. for 2 weeks	18	↔ ^f	↔ ^f	↔ ^f
Lopinavir/ritonavir ^e 400 mg/100 mg b.i.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	18	↑30 (↓15 to ↑47)	↑37 (↓20 to ↑55)	↑52 (↓28 to ↑82)
Methadone 70 to 120 mg q.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	19	R-Methadone (active)		
			↓21 ^g (↓30 to ↓12)	↓18 ^g (↓27 to ↓8)	↓11 ^g (↓21 to ↑1)
			S-Methadone (inactive)		

			↓43 ^g (↓49 to ↓37)	↓43 ^g (↓50 to ↓36)	↓41 ^g (↓49 to ↓31)
Nevirapine 200 mg b.i.d. for 2 weeks ^h	1,400 mg b.i.d. for 2 weeks	17	↑25 (↑14 to ↑37)	↑29 (↑19 to ↑40)	↑34 (↑20 to ↑49)
Nevirapine 200 mg b.i.d. for 2 weeks ^h	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	17	↑13 (↑3 to ↑24)	↑14 (↑5 to ↑24)	↑22 (↑9 to ↑35)
Norethindrone ^c 0.5 mg q.d. for 21 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 21 days	25	↓38 (↓32 to ↓44)	↓34 (↓30 to ↓37)	↓26 (↓20 to ↓32)
Phenytoin 300 mg q.d. for 10 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 10 days	14	↓20 (↓12 to ↓27)	↓22 (↓17 to ↓27)	↓29 (↓23 to ↓34)
Rifabutin 150 mg every other day for 2 weeks ⁱ (25-O-desacetylriofabutin metabolite) Rifabutin + 25-O- desacetylriofabutin metabolite	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	15	↓14 (↓28 to ↑4) ↑579 (↑479 to ↑698) NA	↔ ↑1,120 (↑965 to ↑1,300) ↑64 (↑46 to ↑84)	↑28 (↑12 to ↑46) ↑2,510 (↑1,910 to ↑3,300) NA

^a Concomitant medication is also shown in this column where appropriate.

^b Comparison arm of atazanavir 300 mg q.d. plus ritonavir 100 mg q.d. for 10 days.

^c Administered as a combination oral contraceptive tablet: ethinyl estradiol 0.035 mg/norethindrone 0.5 mg.

^d Patients were receiving LEXIVA/ritonavir for 10 days prior to the 4-day treatment period with both ketoconazole and LEXIVA/ritonavir.

^e Data represent lopinavir concentrations.

^f Compared with lopinavir 400 mg/ritonavir 100 mg b.i.d. for 2 weeks.

^g Dose normalized to methadone 100 mg. The unbound concentration of the active moiety, R-methadone, was unchanged.

^h Patients were receiving nevirapine for at least 12 weeks prior to study.

ⁱ Comparison arm of rifabutin 300 mg q.d. for 2 weeks. AUC is AUC_(0-48 hr).

↑ = Increase; ↓ = Decrease; ↔ = No change (↑ or ↓ less than 10%); ND = Interaction cannot be determined as C_{min} was below the lower limit of quantitation.

Table 13. Drug Interactions: Pharmacokinetic Parameters for Coadministered Drug in the Presence of Amprenavir After Administration of AGENERASE

Coadministered Drug(s) and Dose(s)	Dose of AGENERASE	n	% Change in Pharmacokinetic Parameters of Coadministered Drug (90% CI)		
			C _{max}	AUC	C _{min}
Abacavir 300 mg b.i.d. for 2 to 3 weeks	900 mg b.i.d. for 2 to 3 weeks	4	↔ ^a	↔ ^a	↔ ^a
Clarithromycin 500 mg b.i.d. for 4 days	1,200 mg b.i.d. for 4 days	12	↓10 (↓24 to ↑7)	↔	↔
Delavirdine 600 mg b.i.d. for 10 days	600 mg b.i.d. for 10 days	9	↓47 ^b	↓61 ^b	↓88 ^b
Ethinyl estradiol 0.035 mg for 1 cycle	1,200 mg b.i.d. for 28 days	10	↔	↔	↑32 (↓3 to ↑79)
Indinavir 800 mg t.i.d. for 2 weeks (fasted)	750 mg or 800 mg t.i.d. for 2 weeks (fasted)	9	↓22 ^a	↓38 ^a	↓27 ^a
Ketoconazole 400 mg single dose	1,200 mg single dose	12	↑19 (↑8 to ↑33)	↑44 (↑31 to ↑59)	NA
Lamivudine 150 mg single dose	600 mg single dose	11	↔	↔	NA
Methadone 44 to 100 mg q.d. for >30 days	1,200 mg b.i.d. for 10 days	16	R-Methadone (active)		
			↓25 (↓32 to ↓18)	↓13 (↓21 to ↓5)	↓21 (↓32 to ↓9)
			S-Methadone (inactive)		
			↓48 (↓55 to ↓40)	↓40 (↓46 to ↓32)	↓53 (↓60 to ↓43)
Nelfinavir 750 mg t.i.d. for 2 weeks (fed)	750 mg or 800 mg t.i.d. for 2 weeks (fed)	6	↑12 ^a	↑15 ^a	↑14 ^a
Norethindrone 1 mg for 1 cycle	1,200 mg b.i.d. for 28 days	10	↔	↑18 (↑1 to ↑38)	↑45 (↑13 to ↑88)
Rifabutin 300 mg q.d. for 10 days	1,200 mg b.i.d. for 10 days	5	↑119 (↑82 to ↑164)	↑193 (↑156 to ↑235)	↑271 (↑171 to ↑409)
Rifampin 300 mg q.d. for 4 days	1,200 mg b.i.d. for 4 days	11	↔	↔	ND
Saquinavir 800 mg t.i.d. for 2 weeks (fed)	750 mg or 800 mg t.i.d. for 2 weeks (fed)	7	↑21 ^a	↓19 ^a	↓48 ^a

Zidovudine 300 mg single dose	600 mg single dose	12	↑40 (↑14 to ↑71)	↑31 (↑19 to ↑45)	NA
----------------------------------	-----------------------	----	---------------------	---------------------	----

^a Compared with historical data.

^b Median percent change; confidence interval not reported.

↑ = Increase; ↓ = Decrease; ↔ = No change (↑ or ↓ less than 10%); NA = C_{min} not calculated for single-dose study; ND = Interaction cannot be determined as C_{min} was below the lower limit of quantitation.

12.4 Microbiology

Mechanism of Action: Fosamprenavir is a prodrug that is rapidly hydrolyzed to amprenavir by cellular phosphatases in the gut epithelium as it is absorbed. Amprenavir is an inhibitor of HIV-1 protease. Amprenavir binds to the active site of HIV-1 protease and thereby prevents the processing of viral Gag and Gag-Pol polyprotein precursors, resulting in the formation of immature non-infectious viral particles.

Antiviral Activity: Fosamprenavir has little or no antiviral activity in vitro. The in vitro antiviral activity of amprenavir was evaluated against HIV-1 IIB in both acutely and chronically infected lymphoblastic cell lines (MT-4, CEM-CCRF, H9) and in peripheral blood lymphocytes. The 50% effective concentration (EC₅₀) of amprenavir ranged from 0.012 to 0.08 μM in acutely infected cells and was 0.41 μM in chronically infected cells (1 μM = 0.50 mcg/mL). The median EC₅₀ value of amprenavir against HIV-1 isolates from clades A to G was 0.00095 μM in peripheral blood mononuclear cells (PBMCs). Similarly, the EC₅₀ values for amprenavir against monocytes/macrophage tropic HIV-1 isolates (clade B) ranged from 0.003 to 0.075 μM in monocyte/macrophage cultures. The EC₅₀ values of amprenavir against HIV-2 isolates grown in PBMCs were higher than those for HIV-1 isolates, and ranged from 0.003 to 0.11 μM. Amprenavir exhibited synergistic anti-HIV-1 activity in combination with the nucleoside reverse transcriptase inhibitors (NRTIs) abacavir, didanosine, lamivudine, stavudine, tenofovir, and zidovudine; the non-nucleoside reverse transcriptase inhibitors (NNRTIs) delavirdine and efavirenz; and the protease inhibitors atazanavir and saquinavir. Amprenavir exhibited additive anti-HIV-1 activity in combination with the NNRTI nevirapine, the protease inhibitors indinavir, lopinavir, nelfinavir, and ritonavir; and the fusion inhibitor enfuvirtide. These drug combinations have not been adequately studied in humans.

Resistance: HIV-1 isolates with decreased susceptibility to amprenavir have been selected in vitro and obtained from patients treated with fosamprenavir. Genotypic analysis of isolates from treatment-naïve patients failing amprenavir-containing regimens showed mutations in the HIV-1 protease gene resulting in amino acid substitutions primarily at positions V32I, M46I/L, I47V, I50V, I54L/M, and I84V, as well as mutations in the p7/p1 and p1/p6 Gag and Gag-Pol polyprotein precursor cleavage sites. Some of these amprenavir resistance-associated mutations have also been detected in HIV-1 isolates from antiretroviral-naïve patients treated with LEXIVA. Of the 488 antiretroviral-naïve patients treated with LEXIVA 1,400 mg twice daily or LEXIVA 1,400 mg plus ritonavir 200 mg once daily in studies APV30001 and

APV30002, respectively, 61 patients (29 receiving LEXIVA and 32 receiving LEXIVA/ritonavir) with virologic failure (plasma HIV-1 RNA greater than 1,000 copies/mL on 2 occasions on or after Week 12) were genotyped. Five of the 29 antiretroviral-naïve patients (17%) receiving LEXIVA without ritonavir in study APV30001 had evidence of genotypic resistance to amprenavir: I54L/M (n = 2), I54L + L33F (n = 1), V32I + I47V (n = 1), and M46I + I47V (n = 1). No amprenavir resistance-associated mutations were detected in antiretroviral-naïve patients treated with LEXIVA/ritonavir for 48 weeks in study APV30002. However, the M46I and I50V mutations were detected in isolates from 1 virologic failure patient receiving LEXIVA/ritonavir once daily at Week 160 (HIV-1 RNA greater than 500 copies/mL). Upon retrospective analysis of stored samples using an ultrasensitive assay, these resistant mutants were traced back to Week 84 (76 weeks prior to clinical virologic failure).

Cross-Resistance: Varying degrees of cross-resistance among HIV-1 protease inhibitors have been observed. An association between virologic response at 48 weeks (HIV-1 RNA level less than 400 copies/mL) and protease inhibitor-resistance mutations detected in baseline HIV-1 isolates from protease inhibitor-experienced patients receiving LEXIVA/ritonavir twice daily (n = 88), or lopinavir/ritonavir twice daily (n = 85) in study APV30003 is shown in Table 14. The majority of subjects had previously received either one (47%) or 2 protease inhibitors (36%), most commonly nelfinavir (57%) and indinavir (53%). Out of 102 subjects with baseline phenotypes receiving twice-daily LEXIVA/ritonavir, 54% (n = 55) had resistance to at least one protease inhibitor, with 98% (n = 54) of those having resistance to nelfinavir. Out of 97 subjects with baseline phenotypes in the lopinavir/ritonavir arm, 60% (n = 58) had resistance to at least one protease inhibitor, with 97% (n = 56) of those having resistance to nelfinavir.

Table 14. Responders at Study Week 48 by Presence of Baseline Protease Inhibitor Resistance-Associated Mutations^a

Protease Inhibitor-mutations ^b	LEXIVA/Ritonavir b.i.d. (n = 88)		Lopinavir/Ritonavir b.i.d. (n = 85)	
D30N	21/22	95%	17/19	89%
N88D/S	20/22	91%	12/12	100%
L90M	16/31	52%	17/29	59%
M46I/L	11/22	50%	12/24	50%
V82A/F/T/S	2/9	22%	6/17	35%
I54V	2/11	18%	6/11	55%
I84V	1/6	17%	2/5	40%

^aResults should be interpreted with caution because the subgroups were small.

^bMost patients had greater than 1 protease inhibitor resistance-associated mutation at baseline.

The virologic response based upon baseline phenotype was assessed. Baseline isolates from protease inhibitor-experienced patients responding to LEXIVA/ritonavir twice daily had a median shift in susceptibility to amprenavir relative to a standard wild-type reference strain of 0.7 (range: 0.1 to 5.4, n = 62), and baseline isolates from individuals failing therapy had a median shift in susceptibility of 1.9 (range: 0.2 to 14, n = 29). Because this was a select patient population, these data do not constitute definitive clinical susceptibility break points. Additional data are needed to determine clinically relevant break points for LEXIVA.

Isolates from 15 of the 20 patients receiving twice-daily LEXIVA/ritonavir up to Week 48 and experiencing virologic failure/ongoing replication were subjected to genotypic analysis. The following amprenavir resistance-associated mutations were found either alone or in combination: V32I, M46I/L, I47V, I50V, I54L/M, and I84V. Isolates from 4 of the 16 patients continuing to receive twice-daily LEXIVA/ritonavir up to Week 96 who experienced virologic failure underwent genotypic analysis. Isolates from 2 patients contained amprenavir resistance-associated mutations: V32I, M46I, and I47V in 1 isolate and I84V in the other.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In long-term carcinogenicity studies, fosamprenavir was administered orally for up to 104 weeks at doses of 250, 400, or 600 mg/kg/day in mice and at doses of 300, 825, or 2,250 mg/kg/day in rats. Exposures at these doses were 0.3- to 0.7-fold (mice) and 0.7- to 1.4-fold (rats) those in humans given 1,400 mg twice daily of fosamprenavir alone, and 0.2- to 0.3-fold (mice) and 0.3- to 0.7-fold (rats) those in humans given 1,400 mg once daily of fosamprenavir plus 200 mg ritonavir once daily. Exposures in the carcinogenicity studies were 0.1- to 0.3-fold (mice) and 0.3- to 0.6-fold (rats) those in humans given 700 mg of fosamprenavir plus 100 mg ritonavir twice daily. There was an increase in hepatocellular adenomas and hepatocellular carcinomas at all doses in male mice and at 600 mg/kg/day in female mice, and in hepatocellular adenomas and thyroid follicular cell adenomas at all doses in male rats, and at 835 mg/kg/day and 2,250 mg/kg/day in female rats. The relevance of the hepatocellular findings in the rodents for humans is uncertain. Repeat dose studies with fosamprenavir in rats produced effects consistent with enzyme induction, which predisposes rats, but not humans, to thyroid neoplasms. In addition, in rats only there was an increase in interstitial cell hyperplasia at 825 mg/kg/day and 2,250 mg/kg/day, and an increase in uterine endometrial adenocarcinoma at 2,250 mg/kg/day. The incidence of endometrial findings was slightly increased over concurrent controls, but was within background range for female rats. The relevance of the uterine endometrial adenocarcinoma findings in rats for humans is uncertain.

Fosamprenavir was not mutagenic or genotoxic in a battery of in vitro and in vivo assays. These assays included bacterial reverse mutation (Ames), mouse lymphoma, rat micronucleus, and chromosome aberrations in human lymphocytes.

The effects of fosamprenavir on fertility and general reproductive performance were investigated in male (treated for 4 weeks before mating) and female rats (treated for 2 weeks before mating through postpartum day 6). Systemic exposures ($AUC_{0-24 \text{ hr}}$) to amprenavir in these studies were 3 (males) to 4 (females) times higher than exposures in humans following administration of the MRHD of fosamprenavir alone or similar to those seen in humans following administration of fosamprenavir in combination with ritonavir. Fosamprenavir did not impair mating or fertility of male or female rats and did not affect the development and maturation of sperm from treated rats.

14 CLINICAL STUDIES

14.1 Therapy-Naive Adult Patients

Study APV30001: APV30001 was a randomized, open-label study, comparing treatment with LEXIVA Tablets (1,400 mg twice daily) versus nelfinavir (1,250 mg twice daily) in 249 antiretroviral treatment-naive patients. Both groups of patients also received abacavir (300 mg twice daily) and lamivudine (150 mg twice daily).

The mean age of the patients in this study was 37 years (range: 17 to 70 years), 69% of the patients were males, 20% were CDC Class C (AIDS), 24% were Caucasian, 32% were black, and 44% were Hispanic. At baseline, the median CD4+ cell count was 212 cells/mm³ (range: 2 to 1,136 cells/mm³; 18% of patients had a CD4+ cell count of less than 50 cells/mm³ and 30% were in the range of 50 to less than 200 cells/mm³). Baseline median HIV-1 RNA was 4.83 log₁₀ copies/mL (range: 1.69 to 7.41 log₁₀ copies/mL; 45% of patients had greater than 100,000 copies/mL).

The outcomes of randomized treatment are provided in Table 15.

Table 15. Outcomes of Randomized Treatment Through Week 48 (APV30001)

Outcome (Rebound or discontinuation = failure)	LEXIVA 1,400 mg b.i.d. (n = 166)	Nelfinavir 1,250 mg b.i.d. (n = 83)
Responder ^a	66% (57%)	52% (42%)
Virologic failure	19%	32%
Rebound	16%	19%
Never suppressed through Week 48	3%	13%
Clinical progression	1%	1%
Death	0%	1%
Discontinued due to adverse reactions	4%	2%
Discontinued due to other reasons ^b	10%	10%

^a Patients achieved and maintained confirmed HIV-1 RNA less than 400 copies/mL (less than 50 copies/mL) through Week 48 (Roche AMPLICOR HIV-1 MONITOR Assay Version 1.5).

^b Includes consent withdrawn, lost to follow up, protocol violations, those with missing data, and other reasons.

Treatment response by viral load strata is shown in Table 16.

Table 16. Proportions of Responders Through Week 48 by Screening Viral Load (APV30001)

Screening Viral Load HIV-1 RNA (copies/mL)	LEXIVA 1,400 mg b.i.d.		Nelfinavir 1,250 mg b.i.d.	
	<400 copies/mL	n	<400 copies/mL	n
≤100,000	65%	93	65%	46
>100,000	67%	73	36%	37

Through 48 weeks of therapy, the median increases from baseline in CD4+ cell counts were 201 cells/mm³ in the group receiving LEXIVA and 216 cells/mm³ in the nelfinavir group.

Study APV30002: APV30002 was a randomized, open-label study, comparing treatment with LEXIVA Tablets (1,400 mg once daily) plus ritonavir (200 mg once daily) versus nelfinavir (1,250 mg twice daily) in 649 treatment-naïve patients. Both treatment groups also received abacavir (300 mg twice daily) and lamivudine (150 mg twice daily).

The mean age of the patients in this study was 37 years (range: 18 to 69 years), 73% of the patients were males, 22% were CDC Class C, 53% were Caucasian, 36% were black, and 8% were Hispanic. At baseline, the median CD4+ cell count was 170 cells/mm³ (range: 1 to 1,055 cells/mm³; 20% of patients had a CD4+ cell count of less than 50 cells/mm³ and 35% were in the range of 50 to less than 200 cells/mm³). Baseline median HIV-1 RNA was 4.81 log₁₀ copies/mL (range: 2.65 to 7.29 log₁₀ copies/mL; 43% of patients had greater than 100,000 copies/mL).

The outcomes of randomized treatment are provided in Table 17.

Table 17. Outcomes of Randomized Treatment Through Week 48 (APV30002)

Outcome (Rebound or discontinuation = failure)	LEXIVA 1,400 mg q.d./ Ritonavir 200 mg q.d. (n = 322)	Nelfinavir 1,250 mg b.i.d. (n = 327)
Responder ^a	69% (58%)	68% (55%)
Virologic failure	6%	16%
Rebound	5%	8%
Never suppressed through Week 48	1%	8%
Death	1%	0%
Discontinued due to adverse reactions	9%	6%
Discontinued due to other reasons ^b	15%	10%

^a Patients achieved and maintained confirmed HIV-1 RNA less than 400 copies/mL (less than 50 copies/mL) through Week 48 (Roche AMPLICOR HIV-1 MONITOR Assay Version 1.5).

^b Includes consent withdrawn, lost to follow up, protocol violations, those with missing data, and other reasons.

Treatment response by viral load strata is shown in Table 18.

Table 18. Proportions of Responders Through Week 48 by Screening Viral Load (APV30002)

Screening Viral Load HIV-1 RNA (copies/mL)	LEXIVA 1,400 mg q.d./ Ritonavir 200 mg q.d.		Nelfinavir 1,250 mg b.i.d.	
	<400 copies/mL	n	<400 copies/mL	n
≤100,000	72%	197	73%	194
>100,000	66%	125	64%	133

Through 48 weeks of therapy, the median increases from baseline in CD4+ cell counts were 203 cells/mm³ in the group receiving LEXIVA and 207 cells/mm³ in the nelfinavir group.

14.2 Protease Inhibitor-Experienced Adult Patients

Study APV30003: APV30003 was a randomized, open-label, multicenter study comparing 2 different regimens of LEXIVA plus ritonavir (LEXIVA Tablets 700 mg twice daily plus ritonavir 100 mg twice daily or LEXIVA Tablets 1,400 mg once daily plus ritonavir 200 mg once daily) versus lopinavir/ritonavir (400 mg/100 mg twice daily) in 315 patients who had experienced virologic failure to 1 or 2 prior protease inhibitor-containing regimens.

The mean age of the patients in this study was 42 years (range: 24 to 72 years), 85% were male, 33% were CDC Class C, 67% were Caucasian, 24% were black, and 9% were Hispanic. The median CD4+ cell count at baseline was 263 cells/mm³ (range: 2 to 1,171 cells/mm³).

Baseline median plasma HIV-1 RNA level was 4.14 log₁₀ copies/mL (range: 1.69 to 6.41 log₁₀ copies/mL).

The median durations of prior exposure to NRTIs were 257 weeks for patients receiving LEXIVA/ritonavir twice daily (79% had greater than or equal to 3 prior NRTIs) and 210 weeks for patients receiving lopinavir/ritonavir (64% had greater than or equal to 3 prior NRTIs). The median durations of prior exposure to protease inhibitors were 149 weeks for patients receiving LEXIVA/ritonavir twice daily (49% received greater than or equal to 2 prior protease inhibitors) and 130 weeks for patients receiving lopinavir/ritonavir (40% received greater than or equal to 2 prior protease inhibitors).

The time-averaged changes in plasma HIV-1 RNA from baseline (AAUCMB) at 48 weeks (the endpoint on which the study was powered) were -1.4 log₁₀ copies/mL for twice-daily LEXIVA/ritonavir and -1.67 log₁₀ copies/mL for the lopinavir/ritonavir group.

The proportions of patients who achieved and maintained confirmed HIV-1 RNA less than 400 copies/mL (secondary efficacy endpoint) were 58% with twice-daily LEXIVA/ritonavir and 61% with lopinavir/ritonavir (95% CI for the difference: -16.6, 10.1). The proportions of patients with HIV-1 RNA less than 50 copies/mL with twice-daily LEXIVA/ritonavir and with lopinavir/ritonavir were 46% and 50%, respectively (95% CI for the difference: -18.3, 8.9). The proportions of patients who were virologic failures were 29% with twice-daily LEXIVA/ritonavir and 27% with lopinavir/ritonavir.

The frequency of discontinuations due to adverse events and other reasons, and deaths were similar between treatment arms.

Through 48 weeks of therapy, the median increases from baseline in CD4+ cell counts were 81 cells/mm³ with twice-daily LEXIVA/ritonavir and 91 cells/mm³ with lopinavir/ritonavir.

This study was not large enough to reach a definitive conclusion that LEXIVA/ritonavir and lopinavir/ritonavir are clinically equivalent.

Once-daily administration of LEXIVA plus ritonavir is not recommended for protease inhibitor-experienced patients. Through Week 48, 50% and 37% of patients receiving LEXIVA 1,400 mg plus ritonavir 200 mg once daily had plasma HIV-1 RNA less than 400 copies/mL and less than 50 copies/mL, respectively.

14.3 Pediatric Patients

Two open-label studies in pediatric patients aged 2 to 18 years were conducted. In one study, twice-daily dosing regimens (LEXIVA with or without ritonavir) were evaluated in combination with other antiretroviral agents. A second study evaluated once-daily dosing of LEXIVA with ritonavir; the data from this study were insufficient to support a once-daily dosing regimen in any pediatric patient population.

LEXIVA: Eighteen (16 therapy-naive and 2 therapy-experienced) pediatric patients received LEXIVA Oral Suspension without ritonavir twice daily. At Week 24, 67% (12/18) achieved HIV-1 RNA less than 400 copies/mL, and the median increase from baseline in CD4+ cell count was 353 cells/mm³.

LEXIVA plus ritonavir: Twenty-seven protease inhibitor-naive and 30 protease inhibitor-experienced pediatric patients received LEXIVA Oral Suspension or Tablets with ritonavir twice daily. At Week 24, 70% of protease inhibitor-naive (19/27) and 57% of protease inhibitor-experienced (17/30) patients achieved HIV-1 RNA less than 400 copies/mL; median increases from baseline in CD4+ cell counts were 131 cells/mm³ and 149 cells/mm³ in protease inhibitor-naive and experienced patients, respectively.

16 HOW SUPPLIED/STORAGE AND HANDLING

LEXIVA Tablets, 700 mg, are pink, film-coated, capsule-shaped, biconvex tablets, with “GX LL7” debossed on one face.

Bottle of 60 with child-resistant closure (NDC 49702-207-18).

Store at controlled room temperature of 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) (see USP Controlled Room Temperature). Keep container tightly closed.

LEXIVA Oral Suspension, a white to off-white grape-bubblegum-peppermint-flavored suspension, contains 50 mg of fosamprenavir as fosamprenavir calcium equivalent to approximately 43 mg of amprenavir in each 1 mL.

Bottle of 225 mL with child-resistant closure (NDC 49702-208-53).

This product does not require reconstitution.

Store at 5° to 30°C (41° to 86°F). Shake vigorously before using. Do not freeze.

17 PATIENT COUNSELING INFORMATION

See FDA-approved Patient Labeling (Patient Information).

17.1 Drug Interactions

A statement to patients and healthcare providers is included on the product's bottle label: **ALERT: Find out about medicines that should NOT be taken with LEXIVA.**

LEXIVA may interact with many drugs; therefore, patients should be advised to report to their healthcare provider the use of any other prescription or nonprescription medication or herbal products, particularly St. John's wort.

Patients receiving PDE5 inhibitors should be advised that they may be at an increased risk of PDE5 inhibitor-associated adverse events, including hypotension, visual changes, and priapism, and should promptly report any symptoms to their healthcare provider.

Patients receiving hormonal contraceptives should be instructed to use alternate contraceptive measures during therapy with LEXIVA because hormonal levels may be altered, and if used in combination with LEXIVA and ritonavir, liver enzyme elevations may occur.

17.2 Sulfa Allergy

Patients should inform their healthcare provider if they have a sulfa allergy. The potential for cross-sensitivity between drugs in the sulfonamide class and fosamprenavir is unknown.

17.3 Redistribution/Accumulation of Body Fat

Patients should be informed that redistribution or accumulation of body fat may occur in patients receiving antiretroviral therapy, including LEXIVA, and that the cause and long-term health effects of these conditions are not known at this time.

17.4 Information About Therapy With LEXIVA

Patients should be informed that LEXIVA is not a cure for HIV infection and that they may continue to develop opportunistic infections and other complications associated with HIV disease. The long-term effects of LEXIVA are unknown at this time. Patients should be told that there are currently no data demonstrating that therapy with LEXIVA can reduce the risk of transmitting HIV to others.

Patients should be told that sustained decreases in plasma HIV-1 RNA have been associated with a reduced risk of progression to AIDS and death. Patients should remain under the care of a physician while using LEXIVA. Patients should be advised to take LEXIVA every day as prescribed. LEXIVA must always be used in combination with other antiretroviral drugs. Patients should not alter the dose or discontinue therapy without consulting their physician. If a dose is missed, patients should take the dose as soon as possible and then return to their normal schedule. However, if a dose is skipped, the patient should not double the next dose.

17.5 Oral Suspension

Patients should be instructed to shake the bottle vigorously before each use and that refrigeration of the oral suspension may improve the taste for some patients.

LEXIVA and AGENERASE are registered trademarks of ViiV Healthcare.

The brands listed are trademarks of their respective owners and are not trademarks of ViiV Healthcare. The makers of these brands are not affiliated with and do not endorse ViiV Healthcare or its products.

Manufactured for:



ViiV Healthcare

Research Triangle Park, NC 27709



Vertex Pharmaceuticals Incorporated
Cambridge, MA 02139

by:



GlaxoSmithKline

GlaxoSmithKline

Research Triangle Park, NC 27709

©2011, ViiV Healthcare. All rights reserved.

May 2011

LXV:13PI

PATIENT INFORMATION

LEXIVA[®]
(lex-EE-vah)
(fosamprenavir calcium)
Tablets and Oral Suspension

Read the Patient Information that comes with LEXIVA before you start taking it and each time you get a refill. There may be new information. This information does not take the place of talking with your healthcare provider about your medical condition or treatment. It is important to remain under a healthcare provider's care while taking LEXIVA. Do not change or stop treatment without first talking with your healthcare provider. Talk to your healthcare provider or pharmacist if you have any questions about LEXIVA.

What is the most important information I should know about LEXIVA?

LEXIVA can cause dangerous and life-threatening interactions if taken with certain other medicines. Tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements.

- Some medicines cannot be taken at all with LEXIVA.
- Some medicines will require dose changes if taken with LEXIVA.
- Some medicines will require close monitoring if you take them with LEXIVA.

Know all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements. Keep a list of the medicines you take. Show this list to all your healthcare providers and pharmacists anytime you get a new medicine or refill. Your healthcare providers and pharmacists must know all the medicines you take. They will tell you if you can take other medicines with LEXIVA. Do not start any new medicines while you are taking LEXIVA without talking with your healthcare provider or pharmacist. You can ask your healthcare provider or pharmacist for a list of medicines that can interact with LEXIVA.

What is LEXIVA?

LEXIVA is a medicine you take by mouth to treat HIV infection. HIV is the virus that causes AIDS (acquired immune deficiency syndrome). LEXIVA belongs to a class of anti-HIV medicines called protease inhibitors. LEXIVA is always used with other anti-HIV medicines. When used in combination therapy, LEXIVA may help lower the amount of HIV found in your

blood, raise CD4+ (T) cell counts, and keep your immune system as healthy as possible, so it can help fight infection. However, LEXIVA does not work in all patients with HIV.

LEXIVA does not:

- cure HIV infection or AIDS. We do not know if LEXIVA will help you live longer or have fewer of the medical problems (opportunistic infections) that people get with HIV or AIDS. Opportunistic infections are infections that develop because the immune system is weak. Some of these conditions are pneumonia, herpes virus infections, and *Mycobacterium avium* complex (MAC) infections. It is very important that you see your healthcare provider regularly while you are taking LEXIVA. The long-term effects of LEXIVA are not known.
- lower the risk of passing HIV to other people through sexual contact, sharing needles, or being exposed to your blood. For your health and the health of others, it is important to always practice safer sex by using a latex or polyurethane condom to lower the chance of sexual contact with semen, vaginal secretions, or blood. Never use or share dirty needles.

LEXIVA has not been fully studied in children younger than 2 years or in adults older than 65.

Who should not take LEXIVA?

Do not take LEXIVA if you:

- are taking certain other medicines. Read the section “What is the most important information I should know about LEXIVA?” Do not take the following medicines* with LEXIVA. You could develop serious or life-threatening problems.
 - HALCION® (triazolam; used for insomnia)
 - Ergot medicines: dihydroergotamine, ergonovine, ergotamine, and methylergonovine such as CAFERGOT®, MIGRANAL®, D.H.E. 45®, ergotrate maleate, METHERGINE®, and others (used for migraine headaches)
 - PROPULSID® (cisapride), used for certain stomach problems
 - VERSED® (midazolam), used for sedation
 - ORAP® (pimozide), used for Tourette’s disorder
 - REVATIO® (sildenafil), used for treatment of pulmonary arterial hypertension
 - UROXATRAL® (alfuzosin), used for benign prostatic hyperplasia (BPH)
- are allergic to LEXIVA or any of its ingredients. The active ingredient is fosamprenavir calcium. See the end of this leaflet for a list of all the ingredients in LEXIVA.
- are allergic to AGENERASE (amprenavir).

You should not take AGENERASE (amprenavir) and LEXIVA at the same time.

There are other medicines you should not take if you are taking LEXIVA and NORVIR® (ritonavir) together. You could develop serious or life-threatening problems. Tell your healthcare

provider about all medicines you are taking before you begin taking LEXIVA and NORVIR (ritonavir) together.

What should I tell my healthcare provider before taking LEXIVA?

Before taking LEXIVA, tell your healthcare provider about all of your medical conditions including if you:

- are pregnant or planning to become pregnant. It is not known if LEXIVA can harm your unborn baby. You and your healthcare provider will need to decide if LEXIVA is right for you. If you use LEXIVA while you are pregnant, talk to your healthcare provider about how you can be on the Antiretroviral Pregnancy Registry.
- are breastfeeding. You should not breastfeed if you are HIV-positive because of the chance of passing the HIV virus to your baby through your milk. Also, it is not known if LEXIVA can pass into your breast milk and if it can harm your baby. If you are a woman who has or will have a baby, talk with your healthcare provider about the best way to feed your baby.
- have liver problems. You may be given a lower dose of LEXIVA or LEXIVA may not be right for you.
- have kidney problems
- have diabetes. You may need dose changes in your insulin or other diabetes medicines.
- have hemophilia
- are allergic to sulfa medicines

Before taking LEXIVA, tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements. LEXIVA can cause dangerous and life-threatening interactions if taken with certain other medicines. You may need dose changes in some of your medicines or closer monitoring with some medicines if you also take LEXIVA (see “What is the most important information I should know about LEXIVA.”). Know all the medicines that you take and keep a list of them with you to show healthcare providers and pharmacists.

Women who use birth control pills should choose a different kind of contraception. The use of LEXIVA with NORVIR (ritonavir) in combination with birth control pills may be harmful to your liver. The use of LEXIVA with or without NORVIR may decrease the effectiveness of birth control pills. Talk to your healthcare provider about choosing an effective contraceptive.

How should I take LEXIVA?

- Take LEXIVA exactly as your healthcare provider prescribed.
- Do not take more or less than your prescribed dose of LEXIVA at any one time. Do not change your dose or stop taking LEXIVA without talking with your healthcare provider.
- You can take LEXIVA Tablets with or without food.
- Adults should take LEXIVA Oral Suspension without food.

- Pediatric patients should take LEXIVA Oral Suspension with food. If vomiting occurs within 30 minutes after dosing, the dose should be repeated.
- Shake LEXIVA Oral Suspension vigorously before each use.
- When your supply of LEXIVA or other anti-HIV medicine starts to run low, get more from your healthcare provider or pharmacy. The amount of HIV virus in your blood may increase if one or more of the medicines are stopped, even for a short time.
- Stay under the care of a healthcare provider while using LEXIVA.
- It is important that you do not miss any doses. If you miss a dose of LEXIVA by more than 4 hours, wait and take the next dose at the regular time. However, if you miss a dose by fewer than 4 hours, take your missed dose right away. Then take your next dose at the regular time.
- If you take too much LEXIVA, call your healthcare provider or poison control center right away.

What should I avoid while taking LEXIVA?

- Do not use certain medicines while you are taking LEXIVA. See “What is the most important information I should know about LEXIVA” and “Who should not take LEXIVA?”
- Do not breastfeed. See “Before taking LEXIVA, tell your healthcare provider”. Talk with your healthcare provider about the best way to feed your baby.
- Avoid doing things that can spread HIV infection since LEXIVA doesn't stop you from passing the HIV infection to others.
- Do not share needles or other injection equipment.
- Do not share personal items that can have blood or body fluids on them, like toothbrushes or razor blades.
- Do not have any kind of sex without protection. Always practice safer sex by using a latex or polyurethane condom to lower the chance of sexual contact with semen, vaginal secretions, or blood.

What are the possible side effects of LEXIVA?

LEXIVA may cause the following side effects:

- skin rash. Skin rashes, some with itching, have happened in patients taking LEXIVA. Swelling of the face, lips, and tongue (angioedema) has also been reported. Tell your healthcare provider if you get a rash or develop facial swelling after starting LEXIVA.
- diabetes and high blood sugar (hyperglycemia). Some patients had diabetes before taking LEXIVA while others did not. Some patients may need changes in their diabetes medicine. Others may need a new diabetes medicine.
- increased bleeding problems in some patients with hemophilia.
- worse liver disease. Patients with liver problems, including hepatitis B or C, are more likely to get worse liver disease when they take anti-HIV medicines like LEXIVA.
- changes in blood tests. Some people have changes in blood tests while taking LEXIVA. These include increases seen in liver function tests and blood fat levels, and decreases in white blood

cells. Your healthcare provider may do regular blood tests to see if LEXIVA is affecting your body.

- changes in body fat. These changes have happened in patients taking antiretroviral medicines like LEXIVA. The changes may include an increased amount of fat in the upper back and neck ("buffalo hump"), breast, and around the trunk. Loss of fat from the legs, arms, and face may also happen. The cause and long-term health effects of these conditions are not known at this time.
- kidney stones have been reported in some patients taking LEXIVA. If you develop signs or symptoms of kidney stones (pain in your side, blood in your urine, pain when you urinate) tell your healthcare provider right away.

Common side effects of LEXIVA are nausea, vomiting, and diarrhea. Tell your healthcare provider about any side effects that bother you or that won't go away.

This list of side effects of LEXIVA is not complete. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store LEXIVA?

- LEXIVA Tablets should be stored at room temperature between 59° and 86°F (15° to 30°C). Keep the container of LEXIVA Tablets tightly closed.
- LEXIVA Oral Suspension may be stored at room temperature or refrigerated. Refrigeration of LEXIVA Oral Suspension may improve taste for some patients. Do not freeze.
- Keep LEXIVA and all medicines out of the reach of children.
- Do not keep medicine that is out of date or that you no longer need. Be sure that if you throw any medicine away, it is out of the reach of children.

General information about LEXIVA

Medicines are sometimes prescribed for conditions that are not mentioned in patient information leaflets. Do not use LEXIVA for a condition for which it was not prescribed. Do not give LEXIVA to other people, even if they have the same symptoms you have. It may harm them.

This leaflet summarizes the most important information about LEXIVA. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about LEXIVA that is written for health professionals. For more information you can call toll-free 877-844-8872 or visit www.LEXIVA.com.

What are the ingredients in LEXIVA?

Tablets:

Active Ingredient: fosamprenavir calcium.

Inactive Ingredients: colloidal silicon dioxide, croscarmellose sodium, magnesium stearate, microcrystalline cellulose, and povidone K30. The tablet film-coating contains the inactive ingredients hypromellose, iron oxide red, titanium dioxide, and triacetin.

LEXIVA Tablets, 700 mg, are pink in color and are capsule-shaped, with the letters “GX LL7” printed on one side of the tablet.



Oral Suspension:

Active Ingredient: fosamprenavir calcium

Inactive ingredients: artificial grape-bubblegum flavor, calcium chloride dihydrate, hypromellose, methylparaben, natural peppermint flavor, polysorbate 80, propylene glycol, propylparaben, purified water, and sucralose.

LEXIVA and AGENERASE are registered trademarks of ViiV Healthcare.

* The brands listed are trademarks of their respective owners and are not trademarks of ViiV Healthcare. The makers of these brands are not affiliated with and do not endorse ViiV Healthcare or its products.

Manufactured for:



ViiV Healthcare

Research Triangle Park, NC 27709



Vertex Pharmaceuticals Incorporated
Cambridge, MA 02139



by:

GlaxoSmithKline

Research Triangle Park, NC 27709

©2011, ViiV Healthcare. All rights reserved.

May 2011

LXV: 11PIL